



1st SEM. 2019/2020

UNIVERSITY OF ESWATINI

FINAL EXAMINATION PAPER

**PROGRAMME: B.Sc. in Agricultural Economics and Agribusiness Management
Year 3**

COURSE CODE: AEM307

TITLE OF PAPER: INTRODUCTION TO ECONOMETRICS

TIME ALLOWED: TWO (2) HOURS

INSTRUCTION: 1. ANSWER ALL QUESTIONS
2. EACH QUESTION CARRIES TWENTY FIVE (25) MARKS

**DO NOT OPEN THIS PAPER UNTIL PERMISSION HAS BEEN GRANTED BY
THE CHIEF INVIGILATOR**

QUESTION 1

1. You have been commissioned to investigate the relationship between the birth weights of newborn females and the number of prenatal visits to a physician or midwife that their mothers made during pregnancy. The dependent variable is $bwght_i$, the birth weight of the i -th newborn female, measured in *grams*. The explanatory variable is $pnvisits_i$, the number of prenatal visits of the i -th newborn's mother during pregnancy, measured in number of visits. The model you propose to estimate is given by the population regression equation:

$$bwght_i = \beta_0 + \beta_1 pnvisits_i + \mu$$

Your research assistant has used 857 sample observations on $bwght_i$ and $pnvisits_i$ to estimate the following OLS sample regression equation, where the figures in parentheses below the coefficient estimates are the *estimated standard errors* of the coefficient estimates:

$$bwght_i = 3199.02 + 14.1219 pnvisits_i + \hat{u}_i \quad (i = 1, \dots, N) \quad N = 857$$

(65.6909) (5.36347) ← (standard errors)

- i. Interpret the estimated coefficient on pnvisits. **[3 Marks]**
- ii. Perform a test of the null hypothesis $H_0: \beta_1=0$ against the alternative hypothesis $H_1: \beta_1 \neq 0$ at the 1% significance level (i.e., for significance level $\alpha = 0.01$). Show how you calculated the test statistic. State the decision rule you use, and the inference you would draw from the test. What would you conclude from the results of the test? **[8 MARKS]**
- iii. Compute the two-sided 95% confidence interval for the intercept coefficient β_0 . Use this two-sided 95% confidence interval for β_0 to test the hypothesis that the mean birth weight of newborn females whose mothers made no prenatal visits to a physician or midwife equals 3,000 grams. State the null hypothesis H_0 and the alternative hypothesis H_1 . State the decision rule you use, and the inference you would draw from the test. **[7 MARKS]**
- iv. Perform a test of the proposition that each additional prenatal visit made by the mother is associated on average with an increase in their newborn females' birth weight of less than 25 grams. Use the 5 percent significance level (i.e., $\alpha = 0.05$). State the null hypothesis H_0 and the alternative hypothesis H_1 . Show how you calculated the test statistic. State the decision rule you use, and the inference you would draw from the test. **[7 MARKS]**

QUESTION 2

- i. Discuss the types of data that are used in economic analysis. What are their key characteristics and differences? (You can use example tables while answering this question). **[8 MARKS]**
- ii. What does R-squared measure? How is it calculated and interpreted? **[6 MARKS]**
- iii. Discuss Causality. What is the causal relationship and how is it different from correlation? **[5 MARKS]**
- iv. What are dummy variables? Discuss briefly the features of the dummy variable regression model. **[6 MARKS]**

QUESTION 3

Consider the earnings model: $Wage_i = \beta_1 + \beta_2 Exper_i + \beta_3 Educ_i + u_i$,
 Where $Wage$ is measured in dollars per hour, $Exper$ is work experience in years, and $Educ$ is the number of years of schooling. The OLS regression results for $N = 100$ males in a given year is shown in Table 1 below:

Table 1. STATA results from OLS estimation of the earnings model

Source	SS	df	MS	Number of obs = 100		
				F(2, 97) = 16.47		
Model	2057.5037	2	1028.75185	Prob > F = 0.0000		
Residual	6059.71269	97	62.4712648	R-squared = 0.2535		
				Adj R-squared = 0.2381		
Total	8117.21639	99	81.9920847	Root MSE = 7.9039		

wage	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
Educ	1.435782	.321546	4.47	0.000	.7976026	2.073962
Exper	.328525	.0658247	4.99	0.000	.1978813	.4591687
_cons	-11.91922	4.750254	-2.51	0.014	-21.34716	-2.491275

- i. The researcher performed a correlation test and received the results presented in Table 2 below:

Table 2. Pairwise correlations

	grade	exper	wage
grade	1.0000		
exper	-0.3665	1.0000	
wage	0.2485	0.3163	1.0000

Use the results in Table 2 to discuss the severity of the multicollinearity and the likely impacts on the OLS results in Table 1.

[10 MARKS]

- ii. Using the results in Table 3 below.

Table 3. STATA results with squared OLS residuals as the dependent variable

Source	SS	df	MS	Number of obs = 100		
				F(5, 94) = 1.97		
Model	498933.661	5	99786.7323	Prob > F = 0.0901		
Residual	4759291.93	94	50630.7652	R-squared = 0.0949		
				Adj R-squared = 0.0467		
Total	5258225.59	99	53113.3898	Root MSE = 225.01		
res2	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
Grade	-7.357599	79.35932	-0.09	0.926	-164.9274	150.2122
Exper	-23.67913	16.87954	-1.40	0.164	-57.19386	9.835591
Grade^2	-1.048003	2.223082	-0.47	0.638	-5.461984	3.365978
Exper^2	.270444	.162453	1.66	0.099	-.0521102	.5929982
Exper*Grade	.5788711	.7165818	0.81	0.421	-.8439188	2.001661
_cons	108.2517	582.867	0.19	0.853	-1049.044	1265.548

Perform White's general test for heteroscedasticity. Be sure to carefully set up the null and alternative hypotheses and draw a conclusion.

[15 MARKS]

QUESTION 4

- i. Discuss models with Binary dependent variables. Explain interpretation of Beta in such a model. **[7 MARKS]**
- ii. Discuss the procedure of the Durbin-Watson test and the Breush-Godfrey test for higher order serial correlation. **[11 MARKS]**
- iii. Discuss the procedures of Breusch-Pagan test for heteroskedasticity. **[7 MARKS]**

Percentage Points of the t-Distribution


TABLE D.2
Percentage points of the t distribution

Example

$$\Pr(t > 2.086) = 0.025$$

$$\Pr(t > 1.725) = 0.05 \quad \text{for } df = 20$$

$$\Pr(|t| > 1.725) = 0.10$$



Pr \ df	0.25 0.50	0.10 0.20	0.05 0.10	0.025 0.05	0.01 0.02	0.005 0.010	0.001 0.002
1	1.000	3.078	6.314	12.706	31.821	63.657	318.31
2	0.816	1.886	2.920	4.303	6.965	9.925	22.327
3	0.765	1.638	2.353	3.182	4.541	5.841	10.214
4	0.741	1.533	2.132	2.776	3.747	4.604	7.173
5	0.727	1.476	2.015	2.571	3.365	4.032	5.893
6	0.718	1.440	1.943	2.447	3.143	3.707	5.208
7	0.711	1.415	1.895	2.365	2.998	3.499	4.785
8	0.706	1.397	1.860	2.306	2.896	3.355	4.501
9	0.703	1.383	1.833	2.262	2.821	3.250	4.297
10	0.700	1.372	1.812	2.228	2.764	3.169	4.144
11	0.697	1.363	1.796	2.201	2.718	3.106	4.025
12	0.695	1.356	1.782	2.179	2.681	3.055	3.930
13	0.694	1.350	1.771	2.160	2.650	3.012	3.852
14	0.692	1.345	1.761	2.145	2.624	2.977	3.787
15	0.691	1.341	1.753	2.131	2.602	2.947	3.733
16	0.690	1.337	1.746	2.120	2.583	2.921	3.686
17	0.689	1.333	1.740	2.110	2.567	2.898	3.646
18	0.688	1.330	1.734	2.101	2.552	2.878	3.610
19	0.688	1.328	1.729	2.093	2.539	2.861	3.579
20	0.687	1.325	1.725	2.086	2.528	2.845	3.552
21	0.686	1.323	1.721	2.080	2.518	2.831	3.527
22	0.686	1.321	1.717	2.074	2.508	2.819	3.505
23	0.685	1.319	1.714	2.069	2.500	2.807	3.485
24	0.685	1.318	1.711	2.064	2.492	2.797	3.467
25	0.684	1.316	1.708	2.060	2.485	2.787	3.450
26	0.684	1.315	1.706	2.056	2.479	2.779	3.435
27	0.684	1.314	1.703	2.052	2.473	2.771	3.421
28	0.683	1.313	1.701	2.048	2.467	2.763	3.408
29	0.683	1.311	1.699	2.045	2.462	2.756	3.396
30	0.683	1.310	1.697	2.042	2.457	2.750	3.385
40	0.681	1.303	1.684	2.021	2.423	2.704	3.307
60	0.679	1.296	1.671	2.000	2.390	2.660	3.232
120	0.677	1.289	1.658	1.980	2.358	2.617	3.160
∞	0.674	1.282	1.645	1.960	2.326	2.576	3.090

Selected Upper Percentage Points of the F-Distribution

TABLE D.3
Upper percentage points of the F distribution (continued)

df for denom- inator N_2	df for numerator N_1												
	Pr	1	2	3	4	5	6	7	8	9	10	11	12
22	.25	1.40	1.48	1.47	1.45	1.44	1.42	1.41	1.40	1.39	1.39	1.38	1.37
	.10	2.95	2.56	2.35	2.22	2.13	2.06	2.01	1.97	1.93	1.90	1.88	1.86
	.05	4.30	3.44	3.05	2.82	2.66	2.55	2.46	2.40	2.34	2.30	2.26	2.23
	.01	7.95	5.72	4.82	4.31	3.99	3.76	3.59	3.45	3.35	3.26	3.18	3.12
24	.25	1.39	1.47	1.46	1.44	1.43	1.41	1.40	1.39	1.38	1.38	1.37	1.36
	.10	2.93	2.54	2.33	2.19	2.10	2.04	1.98	1.94	1.91	1.88	1.85	1.83
	.05	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30	2.25	2.21	2.18
	.01	7.82	5.61	4.72	4.22	3.90	3.67	3.50	3.36	3.26	3.17	3.09	3.03
26	.25	1.38	1.46	1.45	1.44	1.42	1.41	1.39	1.38	1.37	1.37	1.36	1.35
	.10	2.91	2.52	2.31	2.17	2.08	2.01	1.96	1.92	1.88	1.86	1.84	1.81
	.05	4.23	3.37	2.98	2.74	2.59	2.47	2.39	2.32	2.27	2.22	2.18	2.15
	.01	7.72	5.53	4.64	4.14	3.82	3.59	3.42	3.29	3.18	3.09	3.02	2.96
28	.25	1.38	1.46	1.45	1.43	1.41	1.40	1.39	1.38	1.37	1.36	1.35	1.34
	.10	2.89	2.50	2.29	2.16	2.06	2.00	1.94	1.90	1.87	1.84	1.81	1.79
	.05	4.20	3.34	2.95	2.71	2.56	2.45	2.36	2.29	2.24	2.19	2.15	2.12
	.01	7.64	5.45	4.57	4.07	3.75	3.53	3.36	3.23	3.12	3.03	2.96	2.90
30	.25	1.38	1.45	1.44	1.42	1.41	1.39	1.38	1.37	1.36	1.35	1.35	1.34
	.10	2.88	2.49	2.28	2.14	2.05	1.98	1.93	1.88	1.85	1.82	1.79	1.77
	.05	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21	2.16	2.13	2.09
	.01	7.56	5.39	4.51	4.02	3.70	3.47	3.30	3.17	3.07	2.98	2.91	2.84
40	.25	1.36	1.44	1.42	1.40	1.39	1.37	1.36	1.35	1.34	1.33	1.32	1.31
	.10	2.84	2.44	2.23	2.09	2.00	1.93	1.87	1.83	1.79	1.76	1.73	1.71
	.05	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12	2.08	2.04	2.00
	.01	7.31	5.18	4.31	3.83	3.51	3.29	3.12	2.99	2.89	2.80	2.73	2.66
60	.25	1.35	1.42	1.41	1.38	1.37	1.35	1.33	1.32	1.31	1.30	1.29	1.29
	.10	2.79	2.39	2.18	2.04	1.95	1.87	1.82	1.77	1.74	1.71	1.68	1.66
	.05	4.00	3.15	2.76	2.53	2.37	2.25	2.17	2.10	2.04	1.99	1.95	1.92
	.01	7.08	4.98	4.13	3.65	3.34	3.12	2.95	2.82	2.72	2.63	2.56	2.50
120	.25	1.34	1.40	1.39	1.37	1.35	1.33	1.31	1.30	1.29	1.28	1.27	1.26
	.10	2.75	2.35	2.13	1.99	1.90	1.82	1.77	1.72	1.68	1.65	1.62	1.60
	.05	3.92	3.07	2.68	2.45	2.29	2.17	2.09	2.02	1.96	1.91	1.87	1.83
	.01	6.85	4.79	3.95	3.48	3.17	2.96	2.79	2.66	2.56	2.47	2.40	2.34
200	.25	1.33	1.39	1.38	1.36	1.34	1.32	1.31	1.29	1.28	1.27	1.26	1.25
	.10	2.73	2.33	2.11	1.97	1.88	1.80	1.75	1.70	1.66	1.63	1.60	1.57
	.05	3.89	3.04	2.65	2.42	2.26	2.14	2.06	1.98	1.93	1.88	1.84	1.80
	.01	6.76	4.71	3.88	3.41	3.11	2.89	2.73	2.60	2.50	2.41	2.34	2.27
∞	.25	1.32	1.39	1.37	1.35	1.33	1.31	1.29	1.28	1.27	1.25	1.24	1.24
	.10	2.71	2.30	2.08	1.94	1.85	1.77	1.72	1.67	1.63	1.60	1.57	1.55
	.05	3.84	3.00	2.60	2.37	2.21	2.10	2.01	1.94	1.88	1.83	1.79	1.75
	.01	6.63	4.61	3.78	3.32	3.02	2.80	2.64	2.51	2.41	2.32	2.25	2.18

Source: Damodar N. Gujarati, *Basic Econometrics*, Third Edition. New York: McGraw-Hill, 1995, p. 814.

TABLE G.4

Critical Values of the Chi-Square Distribution

		Significance Level		
		.10	.05	.01
D e g r e e s o f F r e e d o m	1	2.71	3.84	6.63
	2	4.61	5.99	9.21
	3	6.25	7.81	11.34
	4	7.78	9.49	13.28
	5	9.24	11.07	15.09
	6	10.64	12.59	16.81
	7	12.02	14.07	18.48
	8	13.36	15.51	20.09
	9	14.68	16.92	21.67
	10	15.99	18.31	23.21
	11	17.28	19.68	24.72
	12	18.55	21.03	26.22
	13	19.81	22.36	27.69
	14	21.06	23.68	29.14
	15	22.31	25.00	30.58
	16	23.54	26.30	32.00
	17	24.77	27.59	33.41
	18	25.99	28.87	34.81
	19	27.20	30.14	36.19
	20	28.41	31.41	37.57
	21	29.62	32.67	38.93
	22	30.81	33.92	40.29
	23	32.01	35.17	41.64
	24	33.20	36.42	42.98
	25	34.38	37.65	44.31
	26	35.56	38.89	45.64
	27	36.74	40.11	46.96
	28	37.92	41.34	48.28
	29	39.09	42.56	49.59
	30	40.26	43.77	50.89

Example: The 5% critical value with $df = 8$ is 15.51.

Source: This table was generated using the Stata[®] function