UNIVERSITY OF SWAZILAND

FACULTY OF HEALTH SCIENCES

FINAL EXAMINATION PAPER: MAY 2017

TITLE OF PAPER

Emergency Obstetrics and Neonatal Care

COURSE CODE

: MID 504

DURATION

TWO (2) HOURS

TOTAL MARKS

75

:

INSTRUCTIONS:

- 1. ANSWER ALL QUESTIONS
- 2. FIGURES IN BRACKETS INDICATE MARKS ALLOCATED TO EACH OR PART OF A QUESTION
- 3. ANSWER EACH QUESTION ON A NEW PAGE (ANSWER SHEET)
- 4. QUESTIONS <u>DO NOT</u> CARRY EQUAL MARKS

THIS PAPER SHOULD NOT BE OPENED UNTIL PERMISSION IS GRANTED BY THE INVIGILATOR

Question 1

- 1.1. Differentiate between cord presentation and cord prolapse (4 marks)
- 1. 2. The following are maternal risk factors for cord prolapse: answer with either **True** or **False** (5 marks)
 - a. Prematurity.
 - b. Congenital fetal abnormalities
 - c. Multiple pregnancies.
 - d. Locked twins
 - e. Breech presentation
- 1.3. Explain the immediate management to relieve the pressure on a cord prolapse case (7 marks)
- 1.4. Describe how you would classify hypertension in pregnancy? (4 marks)
- 1.5. The following are <u>NOT</u> signs/symptoms of severe preeclampsia (indicate with **True** or **False**) (5 marks)
 - a. Increased red blood cells
 - b. Visual blurring
 - c. Epigastric pain
 - d. Abnormal liver function
 - e. Elevated platelets

Total Marks = 25

Question 2

- 2.1 How do you classify PPH?(2 marks)
- 2.2 What is the MOST common cause of PPH? (1 mark)
- 2.3 Describe the specific midwifery management for post- partum hemorrhage (7 marks)
- 2.4 The following conditions are risk factors for uterine rupture? State whether **True** or **False** (3 marks)
 - a. Obstructed labour
 - b. Instrumental delivery
 - c. Manipulations, e.g., internal versions
- 2.5. Describe how you would classify uterine inversion. (6 marks)

Total marks = 19

Question 3

- 3.1 Describe how you can differentiate a case of placenta praevia to placenta abruptio, with specific references to the following criteria: abdominal palpation (6 marks), foetal heart conditions (3 marks)? (9 marks)
- 3.2 State the condition that occurs in placenta abruptio when there is extensive infiltration of blood into the myometrium (1 mark)

Total marks= 10

Question 4

A 19 year old primi gravida is admitted at a health centre with history of having delivered at home 7 hours ago. She bled a lot, according to the family, and she became progressively subconscious.

- 4.1 What is your diagnosis on admission?(2 marks)
- 4.2 Describe your actions, as a midwife, on admission. (7 marks)
- 4.3 Based on your response to 4.1, describe the different phases this patient can go through if not well managed. (12 marks)

Total marks= 21

THE END!