

DEPARTMENT OF CHEMISTRY

UNIVERSITY OF SWAZILAND

C204

INTRODUCTION TO ANALYTICAL CHEMISTRY

JULY 2011 SUPPLEMENTARY EXAMINATION

Time Allowed:

Three (3) Hours

Instructions:

1. This examination has six (6) questions. The total number of pages is five (5), including this page.
2. Answer any four (4) questions fully; diagrams should be clear, large and properly labeled. Marks will be deducted for improper units and lack of procedural steps in calculations.
3. Each question is worth 25 marks.

Special Requirements

1. Data sheet.
2. Graph paper.

YOU ARE NOT SUPPOSED TO OPEN THIS PAPER UNTIL PERMISSION TO DO SO HAS BEEN GIVEN BY THE CHIEF INVIGILATOR.

Question 1[25]

- (a) It is found from a reliable method that the glucose level in a blood sample is $125\mu\text{g/g}$. You have developed a new method for glucose analysis, and your values for this blood sample are:

117 $\mu\text{g/g}$ 119 $\mu\text{g/g}$ 111 $\mu\text{g/g}$ 115 $\mu\text{g/g}$ 120 $\mu\text{g/g}$

- (i) Is your method significantly different from the reliable one at 95% confidence level? (2)
- (ii) Is the value 111 $\mu\text{g/g}$ part of the data set? Explain why (2)
- (b) Yet another method has been developed in by another researcher, and the following results are obtained for the sample mentioned above:

135 $\mu\text{g/g}$ 125 $\mu\text{g/g}$ 112 $\mu\text{g/g}$ 122 $\mu\text{g/g}$ 115 $\mu\text{g/g}$ 131 $\mu\text{g/g}$ 119 $\mu\text{g/g}$

- (i) Does your method give the same mean as the one above at 95% confidence level? (2)
- (ii) Does your method give the same precision as the one above at 95% confidence level? (2)
- (c) For the following calibration,
- | | | | | | |
|---|------|------|-------|-------|-------|
| Protein concentration ($\mu\text{g/g}$) | 0.00 | 9.36 | 18.72 | 28.08 | 37.44 |
| Titration volume (mL) | 4.66 | 6.76 | 8.33 | 10.86 | 12.80 |
- (i) Use the least squares method to determine the equation of the best straight line (4)
- (ii) Draw this line (1)
- (d) For diethyl ether with an autoprotolysis constant of 5×10^{-12} as solvent, write down the autoprotolysis equation and calculate the pH of its neutral solutions. (2)
- (e) Calculate the pH of 0.55M NH_4Cl given that K_b for ammonia is 1.76×10^{-5} (3)
- (f) Calculate the fraction dissociated in (e) above (2)
- (g) Explain how you would prepare 500mL of a pH 8.64 buffer solution from solid NH_4Cl and a commercial solution that is 67% v/v NH_3 and density 1.24 g/mL. (5)

Question 2[25]

- (a) (i) Briefly describe how dichlorofluorescein functions as an indicator in the Fajan's precipitation titration. (3)
- (ii) In the determination of chloride ion in waste water, explain why dextrin is added to the solution prior to the Fajans titration. (1)
- (b) Describe the mechanism of formation of a precipitate in gravimetric analysis. (3)
- (c) Using diagrams discuss the two impurity formation process of "occlusion" and "adsorption" in gravimetric analysis. (4)

(d) With regards to the Mohr Method in precipitation titrations:

- (i) Use chemical equations to explain how the end point is detected. (3)
- (ii) Use chemical equations to explain how low pH conditions result in titration errors, and how are these eliminated. (2)
- (iii) Explain why the concentration of indicator is critical in reducing indicator concentration induced errors. (2)

(e) With regards to the Volhard Method in precipitation titrations:

- (i) Use chemical equations to explain how the end point is detected. (3)
- (ii) Use chemical equations to explain why over time the indicator changes colour to the original titrant colour. (2)
- (iii) Describe two ways in which the error resulting in (ii) above is eliminated. (2)

Question 3[25]

(a) For benzoic acid $\text{C}_6\text{H}_5\text{COOH}$ as solvent,

- (i) Write down a chemical equation depicting autoprotolysis (2)
- (ii) Calculate the pH of its "neutral" solutions given that the autoprotolysis constant is 4.5×10^{-6} at 25°C . (3)

(b) In acid-base titrimetry,

- (i) State two reasons why is NaOH not a suitable primary standard (2)
- (ii) Name a common primary standard for the standardization of NaOH (2)

(c) Draw the chemical structure of thymol blue, and explain why this indicator is able to be used over a very wide pH range covering acidic and basic titrations. (4)

(d) Using phenolphthalein as indicator (pink in basic solution but colourless in acid),

- (i) Derive the Henderson – Hasselbach equation for phenolphthalein (3)
- (ii) Given that the K_a of phenolphthalein is 1×10^{-9} use the Henderson equation to derive the useful pH range for the indicator. (4)
- (iii) Plot the titration curve expected from titration of 50ml of 0.02M KOH with 0.100M HBr, using the following data points. (5)

0.00ml added	5.00 ml added	9.99 ml added	10.01 ml added	16.00 ml added
--------------	---------------	---------------	----------------	----------------

Question 4[25]

- (a) In complexometric titrations,
- (i) What does the acronym "DPTA" stand for, and draw its chemical structure (3)
 - (ii) Use an example to explain the term "Multidentate" ligand (3)
 - (iii) For the titration of Mg^{2+} in the presence of Al^{3+} , excess fluoride is usually added at $\text{pH} = 10$. Explain why (3)
- (b) Use an equation to explain what a conditional formation constant is in complexometric titrations. (3)
- (c) Calculate the concentration of free Fe^{3+} in solutions of 0.10M Fe(EDTA)^- at $\text{pH} = 8.00$. (4)
- (d) Draw the chemical structure of Eriochrome Black T, and explain how this indicator works in EDTA titrations. (4)
- (e) A 0.100M EDTA solution is used to titrate 25ml of 0.050M Mg^{2+} buffered at $\text{pH} = 10$ using Eriochrome Black T as indicator.
- (i) Calculate the pMg after addition of the following volumes of EDTA: (4)
- | | | | |
|------|-------|---------|-------|
| 2 mL | 12 mL | 12.5 mL | 13 mL |
|------|-------|---------|-------|
- (ii) Plot the titration curve. (1)

Question 5[25]

- (a) What does the acronym "redox" stand for? (1)
- (b) For the electrochemical cell:
- $$\text{Cd (s)} \mid \text{CdCl}_2(\text{aq}, 0.0538\text{M}) \parallel \text{AgNO}_3(\text{aq}, 0.0328\text{M}) \mid \text{Ag (s)}$$
- (i) What component is represented by the symbol " $\parallel$ "? Explain how it works and why the component is used in electrochemical titrations. (5)
 - (ii) Would the cell be galvanic as written?(5)
- (c) In the iodometric determination of copper using thiosulfate as titrant,
- (i) Name a suitable primary standard (1)
 - (ii) Name a most widely used specific indicator for the titration (1)
 - (iii) Explain the role of ammonium bifluoride added to the samples prior to titration (2)
 - (iv) Explain why the indicator named in c (ii) above is added just before the end point is reached and not at the beginning of the titration (3)
 - (v) Use chemical equations to illustrate how this acts as an indicator for this titration. (3)
- (d) A titration is carried out in a cell, whereby the potential vs SCE (0.241V) is measured for a 25 ml solution of 0.020M Cr^{2+} ($E^0 = -0.41$) titrated with 0.010M Fe^{3+} ($E^0 = 0.770$).
- (i) Calculate the potential (vs SCE) after addition of the following volumes during the titration (3)

5.00ml

50.00ml

100ml

(ii) Sketch the titration curve (1)

Question 6[25]

- (a) Given that at 20°C only 0.24g of an organic acid A dissolves in 100mL of water, but 2.70g of the same acid dissolves in 100mL of ether, calculate the value of the partition coefficient. (4)
- (b) Using diagrams, explain how single stage solvent extraction works. (4)
- (c) List and describe any four (4) properties to be considered in the selection of an organic solvent for extraction. (4)
- (d) List three elements which form stable chloro complexes which are appreciably soluble in organic solvents and used to separate these elements from complex matrices (3)
- (e) Describe two ways of recovering analytes from organic solvent during the stripping stage of solvent extraction. (4)
- (f) Metal chloro complex MCl_3 is extremely soluble in ether, the distribution coefficient for a water/ether system being 50. Calculate the concentration of MCl_3 left in 50ml of aqueous 0.01M FeCl_3 solution after extraction
- (i) once with a 10ml portion of ether (2)
 - (ii) once with a 20ml portion of ether (2)
 - (iii) twice with 10ml portion of ether (2)

1. PERIODIC CHART OF THE ELEMENTS

1 1A H 1.00794	2 2A He 4.00260											13 3A B 10.81	14 4A C 12.011	15 5A N 14.0067	16 6A O 15.9994	17 7A F 18.99840	18 8A Ne 20.179
3 Li 6.941	4 Be 9.01218											5 B 10.81	6 C 12.011	7 N 14.0067	8 O 15.9994	9 F 18.99840	10 Ne 20.179
11 Na 22.98977	12 Mg 24.305	3 3B Al 26.98154	4 4B Si 28.0855	5 5B P 30.97376	6 6B S 32.06	7 7B Cl 35.453	8 8B Ar 39.948										
19 K 39.0983	20 Ca 40.08	21 Sc 44.9559	22 Ti 47.88	23 V 50.9415	24 Cr 51.996	25 Mn 54.9380	26 Fe 55.847	27 Co 58.9332	28 Ni 58.69	29 Cu 63.546	30 Zn 65.38	31 Ga 69.72	32 Ge 72.59	33 As 74.9216	34 Se 78.96	35 Br 79.904	36 Kr 83.80
37 Rb 85.4678	38 Sr 87.62	39 Y 88.9059	40 Zr 91.22	41 Nb 92.9064	42 Mo 95.94	43 Tc (98)	44 Ru 101.07	45 Rh 102.9055	46 Pd 106.42	47 Ag 107.8682	48 Cd 112.41	49 In 114.82	50 Sn 118.69	51 Sb 121.75	52 Te 127.60	53 I 126.9045	54 Xe 131.29
55 Cs 132.9055	56 Ba 137.33	57 La 138.9055	58 Ce 140.12	59 Pr 140.9077	60 Nd 144.24	61 Pm (145)	62 Sm 150.36	63 Eu 151.96	64 Gd 157.25	65 Tb 158.9254	66 Dy 162.50	67 Ho 164.9304	68 Er 167.26	69 Tm 168.9342	70 Yb 173.04	71 Lu 174.967	
87 Fr (223)	88 Ra (226)	89 Ac (227)	90 Th 232.0381	91 Pa 231.0369	92 U 238.0289	93 Np 237.0482	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (260)	

A value in brackets denotes the mass number of the longest lived or best known isotope.

4. NET STABILITY CONSTANTS

Ag(CN) ₂ ⁻	5 × 10
Ag(NH ₃) ₂ ⁺	1.6 × 10
Ag(S ₂ O ₃) ₂ ⁻³	4.7 × 10
Al(OH) ₄ ⁻	1.0 × 10
Ca(EDTA)	1.0 × 10
Cd(CN) ₄	8.3 × 10
Cd(NH ₃) ₄ ⁺⁺	5.5 × 10
Co(NH ₃) ₆ ³⁺	2 × 10
Cr(OH) ₄ ⁻	4 × 10
Cu(CN) ₄ ⁻³	1 × 10
Cu(NH ₃) ₄ ⁺⁺	1.2 × 10
Fe(CN) ₆ ⁻³	4.0 × 10
Fe(CN) ₆ ⁻⁴	2.5 × 10
Fe(SCN) ₄ ⁺⁺	1.0 × 10
HgCl ₄	1.3 × 10
Hg(CN) ₄	8.3 × 10
Hg(SCN) ₄	5.0 × 10
HgI ₄	6.3 × 10
Mg(EDTA)	1.3 × 10
Ni(NH ₃) ₄ ⁺⁺	4.7 × 10
Pb(OH) ₃ ⁻	7.9 × 10
Zn(CN) ₄	4.2 × 10
Zn(NH ₃) ₄ ⁺⁺	7.8 × 10
Zn(OH) ₄	6.3 × 10

2. IONIZATION CONSTANTS (K_a) FOR WEAK ACIDS

Acetic	1.9 × 10 ⁻⁵	Hypochlorous	3.7 × 10 ⁻⁸
2-Amino-pyridinium Ion	2 × 10 ⁻⁷	H ₂ S	K ₁ 9 × 10 ⁻⁸ K ₂ 1 × 10 ⁻¹⁵
Ammonium Ion	5.6 × 10 ⁻¹⁰	Imidazolium Ion	1.1 × 10 ⁻⁷
Anilinium Ion	2.3 × 10 ⁻⁵	Lactic	1.4 × 10 ⁻⁴
Arsenic	K ₁ 5.6 × 10 ⁻³	Methylammonium Ion	2.7 × 10 ⁻¹¹
Benzoic	6.7 × 10 ⁻⁵	Monoethanol-ammonium Ion	3 × 10 ⁻¹⁰
Boric	K ₁ 5 × 10 ⁻¹⁰	Nicotinium Ion	9.6 × 10 ⁻⁹
Carbonic	K ₁ 4.3 × 10 ⁻⁷ K ₂ 5.6 × 10 ⁻¹¹	Oxalic	K ₁ 6 × 10 ⁻² K ₂ 6 × 10 ⁻⁵
Chloroacetic	1.5 × 10 ⁻³	Phenol	1.3 × 10 ⁻¹⁰
Chromic	K ₂ 3.2 × 10 ⁻⁷	Phthalic	K ₁ 4 × 10 ⁻³ K ₂ 4 × 10 ⁻⁶
Citric	K ₁ 8.7 × 10 ⁻⁴ K ₂ 1.8 × 10 ⁻⁵ K ₃ 4 × 10 ⁻⁶	Phosphoric	K ₁ 7.5 × 10 ⁻³ K ₂ 6.2 × 10 ⁻⁸ K ₃ 4.7 × 10 ⁻¹³
Dichloroacetic	5 × 10 ⁻²	Phosphorous	K ₁ 1.0 × 10 ⁻² K ₂ 2.6 × 10 ⁻⁷
EDTA	K ₁ 7 × 10 ⁻³ K ₂ 2 × 10 ⁻³ K ₃ 7 × 10 ⁻⁷ K ₄ 6 × 10 ⁻¹¹	Pyridinium Ion	1 × 10 ⁻⁵
Formic	2 × 10 ⁻⁴	Succinic	K ₁ 7 × 10 ⁻⁵ K ₂ 2.5 × 10 ⁻⁶
α-D(+)-Glucose	5.2 × 10 ⁻¹³	Sulfuric	K ₁ 1.2 × 10 ⁻² K ₂ 6 × 10 ⁻⁸
Glycinium Ion	K ₁ 4.6 × 10 ⁻³ K ₂ 2.5 × 10 ⁻¹⁰	Sulfurous	K ₁ 2 × 10 ⁻² K ₂ 6 × 10 ⁻⁸
Hydrazinium Ion	5.9 × 10 ⁻⁹	Trimethyl-ammonium Ion	1.6 × 10 ⁻¹⁰
Hydrocyanic	7 × 10 ⁻¹⁰	Uric	1.3 × 10 ⁻⁴
Hydrofluoric	7 × 10 ⁻⁴	Water, K _w , 24°C	1.0 × 10 ⁻¹⁴
Hydroxyl-ammonium Ion	9.1 × 10 ⁻⁷		

3. SOLUBILITY PRODUCT CONSTANTS

AgBr	4 × 10 ⁻¹³	BaC ₂ O ₄	2 × 10 ⁻⁸	KClO ₄	2 × 10 ⁻²
Ag ₂ CO ₃	6 × 10 ⁻¹²	BaSO ₄	1 × 10 ⁻¹⁰	MgCO ₃	1 × 10 ⁻⁵
AgCl	1 × 10 ⁻¹⁰	CaCO ₃	5 × 10 ⁻⁹	MgC ₂ O ₄	9 × 10 ⁻⁵
Ag ₂ CrO ₄	2 × 10 ⁻¹²	CaF ₂	4 × 10 ⁻¹¹	MgNH ₄ PO ₄	2 × 10 ⁻¹³
Ag[Ag(CN) ₂]	4 × 10 ⁻¹²	CaC ₂ O ₄	2 × 10 ⁻⁹	Mg(OH) ₂	1 × 10 ⁻¹¹
AgI	1 × 10 ⁻¹⁶	CdS	1 × 10 ⁻²⁸	MnS	1 × 10 ⁻¹⁵
Ag ₃ PO ₄	1 × 10 ⁻¹⁹	Cu(OH) ₂	2 × 10 ⁻²⁰	PbCrO ₄	2 × 10 ⁻¹⁴
Ag ₂ S	1 × 10 ⁻⁵⁰	CuS	1 × 10 ⁻³⁶	PbS	1 × 10 ⁻²⁸
AgCNS	1 × 10 ⁻¹²	Fe(OH) ₃	1 × 10 ⁻³⁶	PbSO ₄	2 × 10 ⁻⁸
Al(OH) ₃	2 × 10 ⁻³²	Hg ₂ Br ₂	3 × 10 ⁻²³	SrCrO ₄	4 × 10 ⁻⁵
BaCO ₃	5 × 10 ⁻⁹	Hg ₂ Cl ₂	6 × 10 ⁻¹⁹	Zn(OH) ₂	3.6 × 10 ⁻¹⁶
BaCrO ₄	1 × 10 ⁻¹⁰	HgS	1 × 10 ⁻⁵²	ZnS	1 × 10 ⁻²⁴

5. FIRST IONIZATION ENERGIES, e.v.

1A	2A	3A	4A	5A	6A	7A
5.4	9.3	8.3	11	13	14	17
5.1	7.6	6.0	8.1	11	10	13
4.3	6.1	6.6	6.8	6.7	7.4	7.9
4.2	5.7	6.6	7.0	6.9	7.2	7.5
3.9	5.2	5.9	5.5	6	6.0	7.0

6. ELECTRONEGATIVITIES, Pauling

1A	2A	3A	4A	5A	6A	7A
1.0	1.5	2.0	2.5	3.0	3.5	4.0
0.9	1.2	1.5	1.8	2.1	2.3	2.6
0.8	1.0	1.3	1.6	1.9	2.2	2.5
0.8	1.0	1.2	1.4	1.6	1.9	2.2
0.7	0.9	1.1	1.3	1.5	1.7	2.0

7. ATOMIC RADII picometers

1A	2A	3A	4A	5A	6A	7A
155	112	98	91	92	73	71
190	160	138	132	128	127	98
235	197	162	147	134	130	126
248	215	178	160	146	139	134
267	222	187	167	149	141	137

8. IONIC RADII pm

Li ⁺	60	Sr ²⁺	113	S ²⁻	184
Na ⁺	95	Ba ²⁺	135	Se ²⁻	198
K ⁺	133	B ³⁺	20	Te ²⁻	221
Rb ⁺	148	Al ³⁺	50	F ⁻	136
Be ²⁺	31	N ³⁺	171	Cl ⁻	181
Mg ²⁺	65	P ³⁺	212	Br ⁻	195
Ca ²⁺	99	O ²⁻	140	I ⁻	216

9. LATTICE ENERGIES

(All negative) kJ/mol	
F ⁻ Cl ⁻ Br ⁻	
Li 1030 840 781	
Na 914 770 728	
K 812 701 671	
Rb 780 682 654	
Cs 744 630 613	

10. HALF LIVES

H ³	12.3 years	K ⁴⁰	1.28 × 10 ⁹ y	I ¹³¹	8.1 day
F ²⁰	11.4 secs	Ca ⁴⁵	165 days	Cs ¹³⁷	30 yea
C ¹⁴	5730 years	Fe ⁵⁹	45 days	Au ¹⁹⁸	2.69 da
Na ²⁴	15.0 hours	Co ⁶⁰	5.26 y	Ra ²²⁶	1620 y
P ³²	14.3 days	Br ⁸²	35.5 hours	U ²³⁵	7.1 × 10
S ³⁵	88 days	Sr ⁹⁰	28 years	U ²³⁸	4.51 × 10
Cl ³⁶	3.1 × 10 ⁵ y	I ¹²⁹	1.7 × 10 ⁷ y	Pu ²³⁹	24,400

Electrode Potentials, E°







Table 26-5 VALUES OF F AT THE 95% CONFIDENCE LEVEL

ν_2	ν_1					
	2	3	4	5	6	∞
2	19.00	19.16	19.25	19.30	19.33	19.50
3	9.55	9.28	9.12	9.01	8.94	8.53
4	6.94	6.59	6.39	6.26	6.16	5.63
5	5.79	5.41	5.19	5.05	4.95	4.36
6	5.14	4.76	4.53	4.39	4.28	3.67
∞	3.00	2.60	2.37	2.21	2.10	1.00

Table 4-2
Values of Student's t

Degrees of freedom	Confidence level ("α")				
	50	80	90	95	99
1	1.000	3.078	6.314	12.706	63.657
2	0.816	1.886	2.920	4.303	9.925
3	0.765	1.638	2.353	3.182	5.841
4	0.741	1.533	2.132	2.776	4.604
5	0.727	1.476	2.015	2.571	4.032
6	0.718	1.440	1.943	2.447	3.707
7	0.711	1.415	1.895	2.365	3.500
8	0.706	1.397	1.860	2.306	3.355
9	0.703	1.383	1.833	2.262	3.250
10	0.700	1.372	1.812	2.228	3.169
15	0.691	1.341	1.753	2.131	2.947
20	0.687	1.325	1.725	2.086	2.845
∞	0.674	1.282	1.645	1.960	2.576

Table 4-4
Values of Q for rejection of data

Q (90% confidence)	0.94	0.76	0.64	0.56	0.51	0.47	0.44	0.41
Number of observations	3	4	5	6	7	8	9	10

Table with 5 columns: Indicator, pH range, pKin, Acid, Base. Rows include Thymol blue, Methyl yellow, Methyl orange, Bromocresol green, Methyl red, Chlorophenol red, Bromothymol blue, Phenol red, Cresol purple, Thymol blue, Phenolphthalein, and Thymolphthalein.

Table with 3 columns: n, Q90, n, Q90, n, Q90. Rows include 3 0.94, 4 0.76, 5 0.64, 6 0.56, 7 0.51, 8 0.47, 9 0.44, 10 0.41. Section 15: Bond Enthalpies. Sub-table with columns: Single, O, N, C, S, F, Cl. Rows include H, C, N, S-S, S-F, S-Cl, H-H, N=N, N=O, C=C, C#C, C=O.

Table with 4 columns: D.F., t90, t90, t95, t9. Rows include 1 1.0, 2 0.82, 3 0.76, 4 0.74, 5 0.73, 6 0.72, 7 0.71, 8 0.71, 9 0.70, 10 0.70, 20 0.69, 30 0.68, infinity 0.67.

12. ELECTRODE POTENTIALS, E°

Na+ + e ⇌ Na -2.713

Mg++ + 2e ⇌ Mg -2.37

Al+++ + 3e ⇌ Al -1.66

Zn++ + 2e ⇌ Zn -0.763

Fe++ + 2e ⇌ Fe -0.44

Cd++ + 2e ⇌ Cd -0.403

Cr+++ + e ⇌ Cr++ -0.38

Tl+ + e ⇌ Tl -0.336

V+++ + e ⇌ V++ -0.255

Sn++ + 2e ⇌ Sn -0.14

Pb++ + 2e ⇌ Pb -0.126

2H+ + 2e ⇌ H2 0.000

SiO2 + 2e ⇌ 2SiO3= 0.09

TiO++ + 2H+ + e ⇌ Ti+++ + H2O 0.10

S + 2H+ + 2e ⇌ H2S 0.14

Sn++ + 2e ⇌ Sn 0.14

Cu++ + e ⇌ Cu+ 0.17

SO4= + 4H+ + 2e ⇌ H2O + H2SO3 0.17

AgCl + e ⇌ Cl- + Ag 0.222

Saturated calomel (0.244)

Hg2Cl2 + 2e ⇌ 2Cl- + 2Hg 0.268

Bi+++ + 3e ⇌ Bi 0.293

UO2++ + 4H+ + 2e ⇌ U++ + 2H2O 0.33

VO++ + 2H+ + e ⇌ V+++ + H2O 0.34

Cu++ + 2e ⇌ Cu 0.34

Fe(CN)6-3 + e ⇌ Fe(CN)6-4 0.355

Cu+ + e ⇌ Cu 0.52

I3- + 2e ⇌ 3I- 0.545

H3AsO4 + 2H+ + 2e ⇌ H3AsO3 + H2O 0.56

I2 + 2e ⇌ 2I- 0.621

2HgCl2 + 2e ⇌ Hg2Cl2 + 2Cl- 0.63

O2 + 2H+ + 2e ⇌ H2O2 0.69

Quinone + 2H+ + 2e ⇌ Hydroquinone 0.70

Fe+++ + e ⇌ Fe++ 0.771

Hg2++ + 2e ⇌ 2Hg 0.792

Ag+ + e ⇌ Ag 0.799

Hg++ + 2e ⇌ Hg 0.851

2Hg++ + 2e ⇌ Hg2++ 0.907

NO3- + 3H+ + 2e ⇌ HNO2 + H2O 0.94

HNO2 + H+ + e ⇌ NO + H2O 0.98

VO2+ + 2H+ + e ⇌ VO++ + H2O 0.999

Br2 + 2e ⇌ 2Br- 1.08

2IO3- + 12H+ + 10e ⇌ 6H2O + I2 1.19

O2 + 4H+ + 4e ⇌ 2H2O 1.229

MnO2 + 4H+ + 2e ⇌ Mn++ + 2H2O 1.23

Cr2O7- + 14H+ + 6e ⇌ 7H2O + 2Cr+++ 1.33

Cl2 + 2e ⇌ 2Cl- 1.358

2BrO3- + 12H+ + 10e ⇌ 6H2O + Br2 1.50

MnO4- + 8H+ + 5e ⇌ 4H2O + Mn++ 1.51

Ce++ + e ⇌ Ce+3 1.61

13. MEAN ACTIVITY COEFFICIENTS

M	KCl	Na2SO4	ZnSO4
0.001	0.965	0.89	0.70
0.01	0.901	0.72	0.39
0.1	0.769	0.45	0.15
1	0.600	0.20	0.045

16. HEATS OF FORMATION

ΔH° in kJ mol-1 at 25°C

All ions in H2O solution except as noted

All Elements = 0

H2	218	H+	0.0	H2Og	-242
O2	249	Na+	-240	H2Ol	-286
Cg	717	Ag+	106	COg	-111
Ng	473	NH4+	-133	CO2g	-394
Fg	79	OH-	-230	NH3g	-46
Clg	122	F-	-333	NOg	90
Brg	112	Cl-	-167	NO2g	33
Ig	107	Br-	-122	N2O4g	9
Sg	279	I-	-55	SO2g	-297
Pg	315	S=	33	SO3g	-396
Na2	107	SO4=	-909	H2Sg	-21
K2	88	CO3=	-677	NaFg	-574
Na+	609	HFg	-271	NaClg	-411
K+	514	HClg	-92	KFg	-567
F-	-255	HBrg	-36	KClg	-437
Cl-	-233	HIg	26	AgClg	-127
CH4g	-75	HCNg	135	AgBrg	-100
C2H2g	227	PH3g	5	PCl3g	-287
C2H4g	52	C6H6g	49	PCl5g	-375
C2H6g	-85	CH3OHl	-238		
C3H8g	-105	C2H5OHl	-235		
nC4H10g	-127	C2H5OHl	-278		
nC5H12g	-209	COCl2g	-219		
CCl4l	-135	CH3Clg	-81		

17. ABS. ENTROPY S°

J mol-1 K-1 at 25°C

H2g	131	P4wh	164	SF6g	292
N2g	192	HFg	174	NOg	211
O2g	205	HClg	187	NO2g	240
Cl2g	223	H2Og	189	N2O4g	304
F2g	203	COg	198	NH3g	192
Cgr	5.7	CO2g	214	PCl3g	312
S8r	254	SO2g	248	PCl5g	365
CH4g	186	SO3g	256	BF3g	254
C2H6g	229	CH3OHl	127		
C3H8g	270	C2H5OHl	283		
C2H2g	201	C2H5OHl	161		
C2H4g	219	(CH3)2Og	266		
C6H6g	269	CH3COOHg	282		

18. ΔG° FORMATION

kJ mol-1 at 25°C

H2	203	HFg	-273	H2Og	-229
F2	62	HClg	-95	H2Ol	-237
Cl2	106	HBrg	-54	SO2g	-300
O2	232	HIg	1.7	SO3g	-371
NOg	87	NH3g	-16	PCl3g	-268
NO2g	51	COg	-137	PCl5g	-305
N2O4g	98	CO2g	-394	CH4g	-51
C2H4g	68	C2H2g	209	C2H6g	-33
C6H6g	125	CH3OHl	-162		
CCl4l	-65	C2H5OHl	-175		
BF3g	-1120	CHCl3g	-70		
SF6g	-1105	CH3COOHg	-374		

20. CONC. ACIDS AND BAS

	M.W.	Density	Wt. %	Mol. it.
Acetic	60.05	1.05	99.5	1
H2SO4	98.07	1.83	94	1
HF	20.01	1.14	45	2
HCl	36.46	1.19	38	1
HBr	80.91	1.52	48	
HNO3	63.01	1.41	69	1
HClO4	100.46	1.67	70	1
H3PO4	98.00	1.69	85	1
NaOH	40.00	1.53	50	1
NH3	17.03	0.90	28	1

21. DENSITIES (g cm-3)

Water at	Air (70 cm)	Mol. it.
0°C 0.9168	Glass	2.7
10° 0.9997	Na2CO3	2.5
20° 0.9982	NaCl	2.2
22° 0.9978	BaSO4	4.5
24° 0.9973	AgCl	5.6
26° 0.9968	Aluminum	2.7
28° 0.9963	Iron	7.9
30° 0.9956	Brass	8.4
90° 0.9653	Mercury	13.6
100° 0.0006	Platinum	21.4

22. MOBILITIES (m2V-1s-1 × 104)

Li+	39	H3O+	350	1/2Ba++
Na+	50	NH4+	73	1/3La+3
K+	74	Ag+	62	1/2SO4=
Cl-	76	OH-	198	1/2PO4-4
Br-	78	I-	77	NO3-

23. WATER V.P. (torr)

°C	4.6	25°
15°	12.8	30°
20°	17.5	50°

24. MISCELLANEOUS

Std. dev. = $\sqrt{\sum (X_i - \bar{X})^2 / (n - 1)}$

Conf. limits = $\bar{X} \pm t_{\alpha} / \sqrt{n}$

$E = E^\circ - (0.0592/n) \log([Red]/[Ox])$

$\log I_1, I_2 = abc = A = \log 1/T$

$\log N_T = \log N_0 - 0.301T/T_1$

$x = (-b \pm \sqrt{b^2 - 4ac}) / 2a$

$n\lambda = 2d \sin \theta$

$2.303 \log_{10} a = \log_e a$

$h = 6.626 \times 10^{-34} \text{ J s}$

$e = 1.602 \times 10^{-19} \text{ C}$

$N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

$F = 96487 \text{ C}$

$g = 9.807 \text{ m s}^{-2}$

$c = 2.998 \times 10^8 \text{ m s}^{-1}$

$1 \text{ amu} = 1.661 \times 10^{-27} \text{ kg}$

$R = 1.987 \text{ cal mol}^{-1} \text{ K}^{-1}$

$= 0.08206 \text{ litre atm mol}^{-1} \text{ K}^{-1}$

$= 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$





$= 8.314 \text{ kPa dm}^3 \text{ mol}^{-1} \text{ K}^{-1}$

$0^\circ \text{C} = 273.15 \text{ K}$

$1 \text{ eV} = 1.602 \times 10^{-19} \text{ J}$

$1 \text{ cal} = 4.1840 \text{ J}$

G / Acid Dissociation Constants

Name	Structure [†]	pK _a [‡]	K _a
Acetic acid (ethanoic acid)	CH ₃ CO ₂ H	4.757	1.75 × 10 ⁻⁵
Alanine	$\begin{array}{c} \text{NH}_3^+ \\ \\ \text{CHCH}_3 \\ \\ \text{CO}_2\text{H} \end{array}$	2.348 (CO ₂ H) 9.857 (NH ₃)	4.49 × 10 ⁻³ 1.36 × 10 ⁻¹⁰
Aminobenzene (aniline)		4.601	2.51 × 10 ⁻⁵
4-Aminobenzenesulfonic acid (sulfanilic acid)		3.232	5.86 × 10 ⁻⁴
2-Aminobenzoic acid (anthranilic acid)		2.08 (CO ₂ H) 4.96 (NH ₃)	8.3 × 10 ⁻³ 1.10 × 10 ⁻⁵
2-Aminoethanethiol (2-mercaptoethylamine)	HSCH ₂ CH ₂ NH ₃ ⁺	8.21 (SH) (μ = 0.1) 10.71 (NH ₃) (μ = 0.1)	6.2 × 10 ⁻⁹ 1.95 × 10 ⁻¹¹
2-Aminoethanol (ethanolamine)	HOCH ₂ CH ₂ NH ₃ ⁺	9.498	3.18 × 10 ⁻¹⁰
2-Aminophenol		4.78 (NH ₃) (20°) 9.97 (OH) (20°)	1.66 × 10 ⁻⁵ 1.05 × 10 ⁻¹⁰
Ammonia	NH ₄ ⁺	9.244	5.70 × 10 ⁻¹⁰
Arginine	$\begin{array}{c} \text{NH}_3^+ \\ \\ \text{CHCH}_2\text{CH}_2\text{CH}_2\text{NHC} \begin{array}{l} \text{NH}_3^+ \\ \text{NH}_2 \end{array} \\ \\ \text{CO}_2\text{H} \end{array}$	1.823 (CO ₂ H) 8.991 (NH ₃) (12.48) (NH ₂)	1.50 × 10 ⁻² 1.02 × 10 ⁻⁹ 3.3 × 10 ⁻¹³
Arsenic acid (hydrogen arsenate)	$\begin{array}{c} \text{O} \\ \\ \text{HO}-\text{As}-\text{OH} \\ \\ \text{OH} \end{array}$	2.24 6.96 11.50	5.8 × 10 ⁻³ 1.10 × 10 ⁻⁷ 3.2 × 10 ⁻¹²

[†] Each acid is written in its protonated form. The acidic protons are indicated in bold type.

[‡] pK_a values refer to 25 °C and zero ionic strength unless otherwise indicated. Values in parentheses are considered to be less reliable. Data are from A. E. Martell and R. M. Smith, *Critical Stability Constants* (New York: Plenum Press, 1974).

N = 14.01 U = 35.45 H = 1.0079

Values of $\alpha_{Y^{4-}}$ for EDTA at 20 C and $\mu = 0.10$ M	
pH	$\alpha_{Y^{4-}}$
0	1.3×10^{-23}
1	1.9×10^{-18}
2	3.3×10^{-14}
3	2.6×10^{-11}
4	3.8×10^{-9}
5	3.7×10^{-7}
6	2.3×10^{-5}
7	5.0×10^{-4}
8	5.6×10^{-3}
9	5.4×10^{-2}
10	0.36
11	0.85
12	0.98
13	1.00
14	1.00

Table 14-2
Formation constants for metal-EDTA complexes

Ion	$\log K_1$	Ion	$\log K_1$	Ion	$\log K_1$
Li ⁺	2.79	Mn ²⁺	25.3 (25 C)	Ce ³⁺	15.98
Na ⁺	1.66	Fe ³⁺	25.1	Pr ³⁺	16.40
K ⁺	0.8	Co ³⁺	41.4 (25 C)	Nd ³⁺	16.61
Rb ⁺	9.2	Zr ⁴⁺	29.5	Pm ³⁺	17.0
Mg ²⁺	8.79	Hf ⁴⁺	29.5 ($\mu = 0.2$)	Sm ³⁺	17.14
Ca ²⁺	10.60	VO ³⁺	18.8	Eu ³⁺	17.35
Sr ²⁺	8.73	VO ₂ ⁺	15.55	Gd ³⁺	17.37
Ba ²⁺	7.86	Ag ⁺	7.32	Tb ³⁺	17.93
Ra ²⁺	7.1	Tl ⁺	6.54	Dy ³⁺	18.30
Sc ³⁺	23.1	Pd ²⁺	18.5 (25 C, $\mu = 0.2$)	Ho ³⁺	18.62
Y ³⁺	18.09			Er ³⁺	18.85
La ³⁺	15.50			Tm ³⁺	19.32
V ²⁺	12.7	Zn ²⁺	16.50	Yb ³⁺	19.51
Cr ²⁺	13.6	Cd ²⁺	16.46	Lu ³⁺	19.83
Mn ²⁺	13.87	Hg ²⁺	21.7	Am ³⁺	17.8 (25 C)
Fe ²⁺	14.32	Sn ²⁺	18.3 ($\mu = 0$)	Cm ³⁺	18.1 (25 C)
Co ²⁺	16.31	Pb ²⁺	18.04	Bk ³⁺	18.5 (25 C)
Ni ²⁺	18.62	Al ³⁺	16.3	Cf ³⁺	18.7 (25 C)
Cu ²⁺	18.80	Ga ³⁺	20.3	Th ⁴⁺	23.2
Ti ³⁺	21.3 (25 C)	In ³⁺	25.0	U ⁴⁺	25.8
V ³⁺	26.0	Tl ³⁺	37.8 ($\mu = 1.0$)	Np ⁴⁺	24.6 (25 C, $\mu = 1.0$)
Cr ³⁺	23.4	Bi ³⁺	27.8		

Note: The stability constant is the equilibrium constant for the reaction $M^{n+} + Y^{4-} \rightleftharpoons MY^{n-4}$. Values in table apply at 20 C, and ionic strength of 1 M, unless otherwise noted.
Source: Data from A. F. Martell and R. M. Smith, *Chemical Stability Constants*, Vol. 1 (New York: Plenum Press, 1974), pp. 204-211.