

UNIVERSITY OF SWAZILAND
BACHELOR OF SCIENCE

SUPPLEMENTARY EXAMINATION 2013

TITLE OF PAPER : PHYSICAL CHEMISTRY

COURSE NUMBER : C202

TIME : 3 HOURS

INSTRUCTIONS : THERE ARE SIX QUESTIONS

: ANSWER ANY FOUR QUESTIONS


: BEGIN THE ANSWER TO EACH QUESTION ON
A SEPARATE SHEET OF PAPER

: DATA SHEETS ARE PROVIDED WITH THIS
EXAMINATION PAPER

DO NOT OPEN THIS PAPER UNTIL THE INVIGILATOR INSTRUCTS YOU TO DO
SO.

Question 1 [25 Marks]

- a) Many gases show nearly ideal behaviour at room temperatures and low pressures. Using a sketch of either an isotherm or compressibility factor 'z' for a real gas and that of an ideal gas, briefly explain how they compare at high densities, moderate densities, and at low densities. [10]
- b) A gaseous mixture in a 250 L container at 125°C contains 16.0 g O₂ and 3.0 g H₂. Assuming ideal gas behaviour calculate:
- partial pressure of each gas [6]
 - total pressure [4]
 - partial pressure of each gas after the mixture has reacted to form water. [5]



Question 2 [25 MARKS]

- a) Using examples and/or diagrams compare and contrast Any Two of the following terms
- reversible and irreversible expansion [5]
 - path and state functions [5]
 - work and heat [5]
 - change in internal energy and change in enthalpy [5]
- b) 2 moles of methane occupies 12 L at 310 K.
- Derive an expression for reversible isothermal expansion. [5]
 - Calculate the work done when the gas expands isothermally against a constant external pressure of 200 torr until its volume has tripled. [5]
 - Calculate the work that would be done if the same expansion in b(ii) occurred in a series of equilibrium steps. [5]

Question 3 [25 Marks]



- a) Write short notes on Any Three of the following concepts:
- Statistical view of entropy [5]
 - Clausius inequality [5]
 - Second law of thermodynamics [5]
 - Third law of thermodynamics [5]

For each concept include the origin or a short derivation showing its origin, an example where applicable and the role or implication of each of the concepts in thermodynamics.

- b) Find $\Delta_f H^\theta$ for the following reactions from standard enthalpies of formation:
- $NH_3(g) + HCl(g) \rightarrow NH_4Cl$ [5]
 - $Cyclopropane(g) \rightarrow propene(g)$ [5]

Question 4 [25 Marks]

- a) (i) Calculate the enthalpy of formation of $\text{N}_2\text{O}_5(\text{g})$ from the following data: [9]




- (ii) Using the enthalpy of formation of $\text{N}_2\text{O}_5(\text{g})$ obtained from a(i) calculate the change in internal energy for the formation of $\text{N}_2\text{O}_5(\text{g})$ [6]

- b) (i) Derive Kirrchoff's equation: [4]

$$\Delta H_r(T_2) = \Delta H_r(T_1) + \Delta_r C_p \Delta T$$

- (ii) Predict the standard enthalpy of reaction at 100°C for the reaction: [6]



Refer to table and the data below:

	$C_p \text{ J mol}^{-1}\text{K}^{-1}$
$\text{N}_2\text{O}_4(\text{g})$	77.28
$\text{NO}_2(\text{g})$	37.20

Question 5 [25 Marks]

- a) Calculate the change in entropies of the system, ΔS_{sys} , the surroundings, ΔS_{surr} , and the total change in entropy, ΔS_{tot} , when a sample of nitrogen gas of mass 14 g at 298 K and 1.00 bar doubles its volume in:

- i) an isothermal reversible expansion [6]

- ii) an irreversible isothermal expansion against an external pressure of 0.5 bar. [4]

- b) What would the change in entropy be if the gas in (a) was compressed to half its volume and simultaneously heated to twice its initial temperature? [5]

- c) If 50g water at 80°C is poured into 100g water at 10°C in an insulated vessel given that $C_{p,m}=75.5 \text{ JK}^{-1}\text{mol}^{-1}$: Calculate:

- i) final temperature of the mixture [4]

- ii) the entropy change [6]


Question 6 [25 Marks]

- a) Derive the integrated Gibbs-Helmholtz equation [5]

$$\frac{\Delta G_2}{T_2} - \frac{\Delta G_1}{T_1} = \Delta H \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

from the fundamental thermodynamic equation $dG = VdP - SdT$

- b) Given the reaction:



Calculate the change in Gibbs free energy ΔG^θ

- i) at 298K [5]
ii) at 500K [5]
iii) Comment on the significance of the values obtained in (i) and (ii). [2]

- c) For the reaction in (b) Calculate

- (i) $\Delta_r H^\theta$ [3]
(ii) ΔU [3]

- (ii). Maximum expansion work, ΔA all at 298 [2]

Useful Relations				General Data	
$(RT)_{298.15K} = 2.4789 \text{ kJ/mol}$				speed of light	$c = 2.997\ 925 \times 10^8 \text{ ms}^{-1}$
$(RT/F)_{298.15K} = 0.025\ 693 \text{ V}$				charge of proton	$e = 1.602\ 19 \times 10^{-19} \text{ C}$
T/K: 100.15 298.15 500.15 1000.15				Faraday constant	$F = Le = 9.648\ 46 \times 10^4 \text{ C mol}^{-1}$
T/Cm ⁻¹ : 69.61 207.22 347.62 695.13				Boltzmann constant	$k = 1.380\ 66 \times 10^{-23} \text{ J K}^{-1}$
1mmHg=133.222 N m ⁻²				Gas constant	$R = Lk = 8.314\ 41 \text{ J K}^{-1} \text{ mol}^{-1}$
$hc/k = 1.438\ 78 \times 10^{-2} \text{ m K}$					$8.205\ 75 \times 10^{-2} \text{ dm}^3 \text{ atm K}^{-1} \text{ mol}^{-1}$
1atm	1 cal	1 eV	1cm ⁻¹		
$-1.01325 \times 10^5 \text{ Nm}^{-2}$	=4.184 J	$=1.602\ 189 \times 10^{-19} \text{ J}$	$=0.124 \times 10^{-3} \text{ eV}$	Planck constant	$h = 6.626\ 18 \times 10^{-34} \text{ Js}$
-760torr		$=96.485 \text{ kJ/mol}$	$=1.9864 \times 10^{-23} \text{ J}$		$\hbar = \frac{h}{2\pi} = 1.054\ 59 \times 10^{-34} \text{ Js}$
-1 bar		$=8065.5 \text{ cm}^{-1}$			
SI-units:				Avogadro constant	$L \text{ or } N_A = 6.022\ 14 \times 10^{23} \text{ mol}^{-1}$
$1 \text{ L} = 1000 \text{ ml} = 1000 \text{ cm}^3 = 1 \text{ dm}^3$				Atomis mass unit	$u = 1.660\ 54 \times 10^{-27} \text{ kg}$
$1 \text{ dm} = 0.1 \text{ m}$				Electron mass	$m_e = 9.109\ 39 \times 10^{-31} \text{ kg}$
$1 \text{ cal (thermochemical)} = 4.184 \text{ J}$				Proton mass	$m_p = 1.672\ 62 \times 10^{-27} \text{ kg}$
dipole moment: 1 Debye = $3.335\ 64 \times 10^{-30} \text{ C m}$				Neutron mass	$m_n = 1.674\ 93 \times 10^{-27} \text{ kg}$
force: $1N = 1 \text{ J m}^{-1} = 1 \text{ kg m s}^{-2} = 10^5 \text{ dyne}$ pressure: $1Pa = 1 \text{ N m}^{-2} = 1 \text{ J m}^{-3}$				Vacuum permittivity	$\epsilon_0 = \mu_0^{-1} c^{-2} = 8.854\ 188 \times 10^{-12} \text{ F}^{-1} \text{ C}^2 \text{ m}^{-1}$
$IJ = 1 \text{ Nm}$ power: $1W = 1 \text{ J s}^{-1}$				Vacuum permeability	$\mu_0 = 4\pi \times 10^{-7} \text{ Js}^2 \text{ C}^{-2} \text{ m}^{-1}$
potential: $1V = 1 \text{ J C}^{-1}$				Bohr magneton	$\mu_B = \frac{e\hbar}{2m_e} = 9.274\ 02 \times 10^{-24} \text{ JT}^{-1}$
magnetic flux: $1T = 1 \text{ V sm}^{-2} = 1 \text{ JCsm}^{-2}$				Nuclear magneton	$\mu_N = \frac{e\hbar}{2m_p} = 5.05079 \times 10^{-27} \text{ JT}^{-1}$
Prefixes:				Gravitational constant	$G = 6.67259 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
p	n	m	m	Gravitational acceleration	$g = 9.80665 \text{ ms}^{-2}$
pico	nano	micro	milli		
10^{-12}	10^{-9}	10^{-6}	10^{-3}	Bohr radius	$a_0 = 5.291\ 77 \times 10^{-11} \text{ m}$
			centi		
			deci		
			kilo		
			mega		
			giga		
			10^{-2}		
			10^{-1}		
			10^3		
			10^6		
			10^9		

THE PERIODIC TABLE OF ELEMENTS

Group	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
	IA	IIA	IIIB	IVB	VIB	VIB	VIB	VIIIB			IB	IIB	IIIA	IVA	VA	VIA	VIIA	VIIIA
Period	1 H 1.008																	
1																		
2	3 Li 6.94	4 Be 9.01																
3	11 Na 22.99	12 Mg 24.31																
4	19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.90	23 V 50.94	24 Cr 52.01	25 Mn 54.9	26 Fe 55.85	27 Co 58.71	28 Ni 58.71	29 Cu 63.54	30 Zn 65.37	31 Ga 69.7	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.91	36 Kr 83.80
5	37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 91.22	42 Mo 95.94	43 Tc 98.9	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3
6	55 Cs 132.9	56 Ba 137.3	71 Lu 174.9	72 Hf 178.5	73 Ta 180.9	74 W 183.8	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 196.9	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 208.9	84 Po 210	85 At 210	86 Rn 222
7	87 Fr 223	88 Ra 226.0	103 Lr 257	104 Unq	105 Unp	106 Unh	107 Uns	108 Uno	109 Une									

Lanthanides		57 La 138.9	58 Ce 140.1	59 Pr 140.9	60 Nd 144.2	61 Pm 146.9	62 Sm 150.9	63 Eu 151.3	64 Gd 157.3	65 Tb 158.9	66 Dy 162.5	67 Ho 164.9	68 Er 167.3	69 Tm 168.9	70 Yb 173.0	
Actinides		89 Ac 227.0	90 Th 232.0	91 Pa 231.0	92 U 238.0	93 Np 237.1	94 Pu 239.1	95 Am 241.1	96 Cm 247.1	97 Bk 249.1	98 Cf 251.1	99 Es 254.1	100 Fm 257.1	101 Md 258.1	102 No 255	

Standard molar Gibbs free energy and molar entropy of formation at 298.15 K

	M _r	ΔG _f ⁰ /KJ/mol	S ⁰ /J K ⁻¹ mol ⁻¹		M _r	ΔG _f ⁰ /KJ/mol	S ⁰ /J K ⁻¹ mol ⁻¹
H ₂ O(g)	18.015	-228.57	188.83	O ₃ (g)	47.998	163.2	238.93
H ₂ O(l)	18.015	-120.35	109.6	NO(g)	30.006	86.55	210.76
H ₂ O ₂ (l)	34.015	-120.35	109.6	NO ₂ (g)	46.006	51.31	240.06
NH ₃ (g)	17.031	-16.45	192.45	N ₂ O ₄ (g)	92.012	97.89	304.29
N ₂ H ₄ (l)	32.045	149.43	121.21	SO ₂ (g)	64.063	-300.19	248.22
N ₃ H(l)	43.028	327.3	140.6	H ₂ S(g)	34.080	-33.56	205.79
N ₃ H(g)	43.028	328.1	238.97	SF ₆ (g)	146.054	-1105.3	291.82
HNO ₃ (l)	63.013	-80.71	155.60	HF(g)	20.006	-273.2	173.78
NH ₂ OH(s)	33.030			HCl(g)	36.461	-95.30	186.91
NH ₄ Cl(s)	53.492	-202.87	94.6	HCl(aq)	36.461	-131.23	56.5
HgCl ₂ (s)	271.50	-178.6	146.0	HBr(g)	80.917	-53.45	198.70
H ₂ SO ₄ (l)	98.078	-690.00	156.90	HI(g)	127.912	1.70	206.59
H ₂ SO ₄ (aq)	98.078	-744.53	20.1	CO ₂ (g)	44.010	-394.36	213.74
NaCl(s)	58.443	-384.14	72.13	CO(g)	28.011	-137.17	197.67
NaOH(s)	39.997	-379.49	64.46	Al ₂ O ₃ (n,s)	101.945	-1582.3	50.92
KCl(s)	74.555	-409.14	82.59	SiO ₂	60.09	-856.64	41.84
KBr(s)	119.011	-380.66	95.90	FeS(s)	87.91	-100.4	60.29
KI(s)	166.006	-324.89	106.32	FeS ₂ (s)	119.975	-166.9	52.93
				AgCl(s)	143.323	-109.79	96.2
He(g)	4.003	0	126.15	Hg(g)	200.59	31.82	174.96
Ar(g)	39.95	0	154.84	Hg(l)	200.59	0	76.02
H ₂ (g)	2.016	0	130.684	Ag(g)	107.87	245.65	173.00
N ₂ (g)	28.013	0	191.61	Ag(s)	107.87	0	42.55
O ₂ (g)	31.999	0	205.138	Na(g)	370.95	76.76	153.71
O ₃ (g)	47.998	163.2	238.93	Na(s)	22.99	0	51.21
Cl ₂ (g)	70.91	0	223.07				
Br ₂ (g)	159.82	3.110	245.46				
Br ₂ (l)	159.82	0	152.23				
I ₂ (g)	253.81	19.33	260.69				
I ₂ (s)	253.81	0	116.135				

organic compounds	M _r	ΔG _f ⁰ /KJ/mol	S ⁰ /J K ⁻¹ mol ⁻¹
CH ₄ (g) methane	16.043	-50.72	186.26
C ₂ H ₂ (g) ethyne	26.038	209.20	200.94
C ₂ H ₄ (g) ethene	28.05	68.15	219.56
C ₂ H ₆ (g) ethane	30.070	-32.82	229.60
C ₃ H ₆ cyclopropane(g)	42.081	104.45	237.55
C ₃ H ₆ propene(g)	42.081	62.78	267.05
C ₄ H ₁₀ n-butane (g)	58.124	-17.03	310.23
C ₅ H ₁₂ n-pentane(g)	72.151	-8.20	348.40
C ₆ H ₁₂ cyclohexane (l)	84.163	26.8	
C ₆ H ₁₄ n-hexane (l)	86.178		204.3
C ₆ H ₆ benzene (l)	78.115	124.3	173.3
C ₆ H ₆ benzene (g)	78.115	129.72	269.31
C ₈ H ₁₈ n-octane (l)	114.233	6.4	361.1
C ₁₀ H ₈ naphthalene (l)	128.175		
CH ₃ OH (g)	32.042	-161.96	239.81
CH ₃ OH (l)	32.042	-166.27	126.8
CH ₃ CHO (g)	44.054	-128.86	250.3
CH ₃ CH ₂ OH (l)	46.07	-174.78	160.7
CH ₃ COOH (l)	60.053	-389.9	159.8
CH ₃ COOC ₂ H ₅ (l)	88.107	-332.7	259.4
C ₆ H ₅ OH (s)	94.114	-50.9	146.0
C ₆ H ₅ NH ₂ (l)	93.129		
CH ₂ (NH ₂)CO ₂ H, glycine (s)	75.068	-373.4	103.5
C ₆ H ₁₂ O ₆ , α-D-glucose (s)	180.159		
C ₆ H ₂₂ O ₁₁ , β-D-glucose (s)	180.159	-910	212
C ₁₂ H ₂₂ O ₁₁ , sucrose (s)	342.303	-1543	360.2
CH ₃ CH(OH)COOH	90.079		
lactic acid (s)			

Source: American Institute of Physics handbook, McGraw-Hill.

Standard molar enthalpies of formation at 298.15 K

	M _r	ΔH _f ^θ /KJ/mol		M _r	ΔH _f ^θ /KJ/mol
H ₂ O(g)	18.015	-241.8	O ₃ (g)	47.998	+142.7
H ₂ O(l)	18.015	-285.8	NO(g)	30.006	+90.2
H ₂ O ₂ (l)	34.015	-187.8	NO ₂ (g)	46.006	+33.2
NH ₃ (g)	17.031	-46.1	N ₂ O ₄ (g)	92.012	+9.2
N ₂ H ₄ (l)	32.045	+50.6	SO ₂ (g)	64.063	-296.8
N ₃ H(l)	43.028	+264.1	H ₂ S(g)	34.080	-20.6
N ₃ H(g)	43.028	+294.1	SF ₆ (g)	146.054	-1209
HNO ₃ (l)	63.013	-174.1	HF(g)	20.006	-271.1
NH ₂ OH(s)	33.030	-114.2	HCl(g)	36.461	-92.3
NH ₄ Cl(s)	53.492	-314.4	HCl(aq)	36.461	-167.2
HgCl ₂ (s)	271.50	-224.3	HBr(g)	80.917	+36.4
H ₂ SO ₄ (l)	98.078	-814.0	HI(g)	127.912	+26.5
H ₂ SO ₄ (aq)	98.078	-909.3	CO ₂ (g)	44.010	-393.5
NaCl(s)	58.443	-411.0	CO(g)	28.011	-110.5
NaOH(s)	39.997	-426.7	Al ₂ O ₃ (α,s)	101.945	-1675.7
KCl(s)	74.555	-435.9	SiO ₂ (s)	60.085	-910.9
KBr(s)	119.011	-392.2	FeS(s)	87.91	-100.0
KI(s)	166.006	-327.6	FeS ₂ (s)	119.975	-178.2
Diatomics(g)	—	0	AgCl(s)	143.323	-127.1

Temperature dependence of heat capacities, C_{p,m} = a+bT+cT⁻²

	a/J K ⁻¹ mol ⁻¹	b/10 ⁻³ J K ⁻² mol ⁻¹	c/10 ⁵ J Kmol ⁻¹
Gases (298-2000K)			
He, Ne, Ar, Kr, Xe	20.78	0	0
H ₂	27.28	3.26	0.50
O ₂	29.96	4.18	-1.67
N ₂	28.56	3.77	-0.50
Cl ₂	37.03	0.67	-2.85
CO ₂	44.23	8.79	-8.62
H ₂ O	30.54	10.29	0
NH ₃	29.75	25.10	-1.55
CH ₄	23.64	47.86	-1.92
Standard molar enthalpies of formation and combustion at 298.15 K.			
	M _r	ΔH _f ^θ /KJ/mol	ΔH _c ^θ /KJ/mol
CH ₄ (g)	16.043	-74.81	
C ₂ H ₂ (g)	26.038	+226.8	1300
C ₂ H ₄ (g)	28.054	+52.30	1411
C ₂ H ₆ (g)	30.070	-84.64	1560
C ₃ H ₆ cyclopropane(g)	42.081	53.35	2091
C ₃ H ₆ (propene(g))	42.081	20.5	2058
C ₄ H ₁₀ n-butane (g)	58.124	-126.11	2877
C ₅ H ₁₂ n-pentane(g)	72.151	-146.4	3536
C ₆ H ₁₂ cyclohexane (l)	84.163	-156.2	3920
C ₆ H ₁₄ n-hexane (l)	86.178	-198.7	4163
C ₆ H ₆ benzene (l)	78.115	+48.99	3268
C ₈ H ₁₈ n-octane (l)	114.233	-249.8	5471
C ₁₀ H ₈ naphthalene (l)	128.175	+78.53	5157
CH ₃ OH (l)	32.042	-239.0	726.1
CH ₃ CHO (g)	44.054	-166.0	1193
CH ₃ CH ₂ OH (l)	46.070	-277.0	1368
CH ₃ COOH (l)	60.053	-484.2	874.5
CH ₃ COOC ₂ H ₅ (l)	88.107	-486.6	2231
C ₆ H ₅ OH (s)	94.114	-165.0	3054
C ₆ H ₅ NH ₂ (l)	93.129	-31.1	3393
NH ₂ CO.NH, urea(s)	60.056	-333.0	632.2
CH ₂ (NH ₂)CO ₂ H, glycine (s)	75.068	-537.2	964.4
C ₆ H ₁₂ O ₆ , α-D-glucose (s)	180.159	-1274	2802
C ₆ H ₂₂ O ₁₁ , β-D-glucose (s)	180.159	-1268	2808
C ₁₂ H ₂₂ O ₁₁ , sucrose (s)	342.303	-2222	5645
CH ₃ CH(OH)COOH	90.079	-694.0	1344
lactic acid (s)			

Standard molar Gibbs free energy and molar entropy of formation at 298.15 K

	M _r	$\Delta G_f^0 / \text{kJ/mol}$	$S^0 / \text{J K}^{-1} \text{ mol}^{-1}$		M _r	$\Delta G_f^0 / \text{kJ/mol}$	$S^0 / \text{J K}^{-1} \text{ mol}^{-1}$
H ₂ O(g)	18.015	-228.57	188.83	O ₃ (g)	47.998	163.2	238.93
H ₂ O(l)	18.015	-120.35	109.6	NO(g)	30.006	86.55	210.76
H ₂ O ₂ (l)	34.015	-120.35	109.6	NO ₂ (g)	46.006	51.31	240.06
NH ₃ (g)	17.031	-16.45	192.45	N ₂ O ₄ (g)	92.012	97.89	304.29
N ₂ H ₄ (l)	32.045	149.43	121.21	SO ₂ (g)	64.063	-300.19	248.22
N ₃ H(l)	43.028	327.3	140.6	H ₂ S(g)	34.080	-33.56	205.79
N ₃ H(g)	43.028	328.1	238.97	SF ₆ (g)	146.054	-1105.3	291.82
HNO ₃ (l)	63.013	-80.71	155.60	HF(g)	20.006	-273.2	173.78
NH ₂ OH(s)	33.030			HCl(g)	36.461	-95.30	186.91
NH ₄ Cl(s)	53.492	-202.87	94.6	HCl(aq)	36.461	-131.23	56.5
HgCl ₂ (s)	271.50	-178.6	146.0	HBr(g)	80.917	-53.45	198.70
H ₂ SO ₄ (l)	98.078	-690.00	156.90	HI(g)	127.912	1.70	206.59
H ₂ SO ₄ (aq)	98.078	-744.53	20.1	CO ₂ (g)	44.010	-394.36	213.74
NaCl(s)	58.443	-384.14	72.13	CO(g)	28.011	-137.17	197.67
NaOH(s)	39.997	-379.49	64.46	Al ₂ O ₃ (α ,s)	101.945	-1582.3	50.92
KCl(s)	74.555	-409.14	82.59	SiO ₂	60.09	-856.64	41.84
KBr(s)	119.011	-380.66	95.90	FeS(s)	87.91	-100.4	60.29
KI(s)	166.006	-324.89	106.32	FeS ₂ (s)	119.975	-166.9	52.93
				AgCl(s)	143.323	-109.79	96.2
He(g)	4.003	0	126.15	Hg(g)	200.59	31.82	174.96
Ar(g)	39.95	0	154.84	Hg(l)	200.59	0	76.02
H ₂ (g)	2.016	0	130.684	Ag(g)	107.87	245.65	173.00
N ₂ (g)	28.013	0	191.61	Ag(s)	107.87	0	42.55
O ₂ (g)	31.999	0	205.138	Na(g)	370.95	76.76	153.71
O ₃ (g)	47.998	163.2	238.93	Na(s)	22.99	0	51.21
Cl ₂ (g)	70.91	0	223.07				
Br ₂ (g)	159.82	3.110	245.46				
Br ₂ (l)	159.82	0	152.23				
I ₂ (g)	253.81	19.33	260.69				
I ₂ (s)	253.81	0	116.135				

	M _r	$\Delta G_f^0 / \text{kJ/mol}$	$S^0 / \text{J K}^{-1} \text{ mol}^{-1}$
organic compounds			
CH ₄ (g) methane	16.043	-50.72	186.26
C ₂ H ₂ (g) ethyne	26.038	209.20	200.94
C ₂ H ₄ (g) ethene	28.05	68.15	219.56
C ₂ H ₆ (g) ethane	30.070	-32.82	229.60
C ₃ H ₆ cyclopropane(g)	42.081	104.45	237.55
C ₃ H ₆ propene(g)	42.081	62.78	267.05
C ₄ H ₁₀ n-butane (g)	58.124	-17.03	310.23
C ₅ H ₁₂ n-pentane(g)	72.151	-8.20	348.40
C ₆ H ₁₂ cyclohexane (l)	84.163	26.8	
C ₆ H ₁₄ n-hexane (l)	86.178		204.3
C ₆ H ₆ benzene (l)	78.115	124.3	173.3
C ₆ H ₆ benzene (g)	78.115	129.72	269.31
C ₈ H ₁₈ n-octane (l)	114.233	6.4	361.1
C ₁₀ H ₈ naphthalene (l)	128.175		
CH ₃ OH (g)	32.042	-161.96	239.81
CH ₃ OH (l)	32.042	-166.27	126.8
CH ₃ CHO (g)	44.054	-128.86	250.3
CH ₃ CH ₂ OH (l)	46.07	-174.78	160.7
CH ₃ COOH (l)	60.053	-389.9	159.8
CH ₃ COOC ₂ H ₅ (l)	88.107	-332.7	259.4
C ₆ H ₅ OH (s)	94.114	-50.9	146.0
C ₆ H ₅ NH ₂ (l)	93.129		
CH ₂ (NH ₂)CO ₂ H, glycine (s)	75.068	-373.4	103.5
C ₆ H ₁₂ O ₆ , α -D-glucose (s)	180.159		
C ₆ H ₂₂ O ₁₁ , β -D-glucose (s)	180.159	-910	212
C ₁₂ H ₂₂ O ₁₁ , sucrose (s)	342.303	-1543	360.2
CH ₃ CH(OH)COOH	90.079		
lactic acid (s)			

Source: American Institute of Physics handbook, McGraw-Hill.

Heat capacities at 25°C

	$C_{v,m}$	$C_{p,m}$
	$\text{JK}^{-1} \text{mol}^{-1}$	$\text{JK}^{-1} \text{mol}^{-1}$
He, Ne, Ar, Kr, Xe	12.47	20.78
H ₂	20.50	28.81
O ₂	21.01	29.33
N ₂	20.83	29.14
CO ₂	28.83	37.14
NH ₃	27.17	35.48
CH ₄	27.43	35.74

F.P Depression, B.P. Elevation

Solvent	F.P °C	K _f °C kg mol ⁻¹	B.P (°C, 101kNm ⁻²)	K _b °C kg mol ⁻¹
Water	0	1.86	100.0	0.52
Benzene	5.51	5.10	80.1	2.60
Acetic Acid	16.6	3.90	118.1	3.10
Cyclohexane	6.5	20.2	81.4	2.79
Camphor	177.7	40.0	205	-
Nitrobenzene	5.7	6.9	210.9	5.24
Ethanol	-177		78.5	1.22
Chloroform	-64		61.3	3.63

Third Law entropies at 25°C, Sm^θ/J K⁻¹ mol⁻¹

Solids	Liquids	Gases
Ag	42.68	H ₂
C(gr)	5.77	Br ₂
C(d)	2.44	O ₂
Cu	33.4	Cl ₂
Zn	41.6	H ₂ O
I ₂	116.7	70.0
S(Rh)	31.9	HNO ₃
AgCl	96.2	C ₂ H ₅ OH
AgBr	104.6	CH ₃ OH
CuSO ₄ ·5H ₂ O	305.4	C ₆ H ₆
HgCl ₂	144	CH ₃ COOH
Sucrose	360.2	C ₆ H ₁₂
		130.6
		192.1
		205.1
		223.0
		213.7
		186.8
		205.6
		192.5
		186.1
		229.4
		265.7