

UNIVERSITY OF SWAZILAND
BACHELOR OF SCIENCE

MAIN EXAMINATION 2018

- TITLE OF PAPER : INTRODUCTION TO THERMODYNAMICS /
(COURSE CODE) (CHE241)
- INTRODUCTORY PHYSICAL CHEMISTRY
(C202)
- TIME : 3 HOURS
- INSTRUCTIONS : THERE ARE SIX QUESTIONS
- : ANSWER ANY FOUR QUESTIONS
- : BEGIN THE ANSWER TO EACH QUESTION ON
A SEPARATE SHEET OF PAPER
- : DATA SHEETS ARE PROVIDED WITH THIS
EXAMINATION PAPER

DO NOT OPEN THIS PAPER UNTIL THE INVIGILATOR INSTRUCTS YOU TO DO SO.

Question 1(25 marks)

- a) (i) Define the variable, compressibility factor, z. [5]
(ii) With the aid of Lennard-Jones potential plot, compressibility and isotherm plots, compare and contrast real and ideal gases.

Your account should make mention of interactions, equations and any necessary theories to help clarify your discussion. [10]

- b) Write short notes on any One of the following:

- i) Virial equation [10]
ii) van der Waal's equation [10]

Use diagrams, equations or plots to clarify your notes where necessary.

QUESTION 2 [25 marks]

- a) A real gas equation of state for a gas is given by:

$$P = RT(V_m - \beta)^{-1} - (\alpha/T)V_m^{-2} \quad (1)$$

- (i) Derive an expression for $V_{m,c}$, T_c and P_c . [12]
(ii) Find an expression for the Boyle's temperature, T_B . [4]
(iii) Estimate the temperature at which oxygen behaves as an ideal gas, T_B given the constants: $\alpha=1.748 \text{ L}^2 \text{ atm mol}^{-2} \text{ K}$ and $\beta= 0.0345 \text{ L mol}^{-1}$. [2]
(iv) Estimate the radii of real gas molecules using equation (1) for real gases given a critical molar volume of $250 \text{ cm}^3 \text{ mol}^{-1}$ [4]

- b) Using the critical point expressions for $V_{m,c}$, T_c and P_c find an expression or value for compressibility at the critical point, Z_c [3]

Question 3 [25 Marks]

- a) Write short notes on the following

- i) Enthalpy change [5]
ii) Hess's Law [5]

- b) Derive Kirchhoff's equation: [6]

$$\Delta H_r(T_2) = \Delta H_r(T_1) + \Delta_r C_{p,m} \Delta T \quad (2)$$

where $C_{p,m}$ is temperature independent.

- b) Using the data in the table below and equation (2) calculate

- i) $\Delta_r H^\theta$ at 298 K [4]
ii) $\Delta_r H$ at 346 K [5]

for the hydrogenation reaction:




Table 1: Molar heat capacities and molar enthalpies of formation

	$\text{C}_2\text{H}_4(\text{g})$	$\text{H}_2(\text{g})$	$\text{C}_2\text{H}_2(\text{g})$
$C_{p,m} \text{ J/mol/K}$	43.56	43.93	28.82
$\Delta_f H^\theta \text{ kJ/mol}$	+52.30	0	+226.8

Question 4 [25 Marks]

- a) Using examples and/or diagrams compare and contrast any one pair of the following terms
- Reversible and irreversible expansion [10]
 - Path and state functions [10]
- b) 4 moles of pentane occupies 25 L at 315 K.
- Derive an expression work for a reversible isothermal expansion process [6]
 - Calculate the work done and heat involved when the gas expands isothermally against a constant external pressure of 115 torr until its volume has doubled. [4]
 - Calculate the efficiency of the system in 1 b (ii) above. [5]

Question 5 [25 Marks]

- a) Write short notes on the following:
- Gibbs Free Energy [4]
 - Helmholtz Function [4]


Use equations, examples and Application to clarify your discussion.

- b) Derive the integrated Gibbs-Helmholtz equation [5]

$$\frac{\Delta G_2}{T_2} - \frac{\Delta G_1}{T_1} = \Delta H \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

starting from the fundamental thermodynamic equation $dG = VdP - SdT$

- c) Given the reaction:



Calculate the change in Gibbs free energy ΔG^θ

- at 298K [5]
- at 500K [5]
- Comment on the significance of the values obtained in (i) and (ii). [2]

Question 6 [25 Marks]

- a) Define internal energy change [10]
- b) To Calibrate a calorimeter a 0.120 g naphthalene, $C_{10}H_8(s)$, was burnt at constant volume and it caused the temperature of the calorimeter to rise by 3.05 K. Then 0.10 g of an unknown compound was burned in the same calorimeter, causing a temperature rise of 2.05 K.
- Calculate the heat capacity of the calorimeter [3]


ii) Is the unknown compound phenol, $C_6H_5OH(s)$ or ethanol, $CH_3CH_2OH(l)$ whose enthalpies of combustion are $\Delta_fH^\theta = -3054 \text{ kJ mol}^{-1}$ and $-1368 \text{ kJ mol}^{-1}$ respectively. [4]

c) Calculate the standard enthalpies of formation of:


i) $KClO_3(s)$ from the enthalpy of formation of KCl [4]

ii) $NOCl(g)$ from the enthalpy of formation of NO [4]

Given the attached table and the following information:



$$\Delta_fH^\theta = -89.4 \text{ kJ/mol}$$



$$\Delta_fH^\theta = +76.5 \text{ kJ/mol}$$

Useful information:

	Molecular weights/g mol ⁻¹
Benzoic acid	122.12
D-ribose $C_5H_{10}O_5(s)$	150.13

THE PERIODIC TABLE OF ELEMENTS

Group	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
	IA	IIA	IIIB	IVB	VB	VIB	VIIB		VIIIB		IB	IIB	IIIA	IVA	VA	VIA	VIIA	VIIIA	
Period 1	1 H 1.008																2 He 4.003		
2	3 Li 6.94	4 Be 9.01												5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18
3	11 Na 22.99	12 Mg 24.31												13 Al 26.9	14 Si 28.09	15 P 30.97	16 S 32.06	17 Cl 35.45	18 Ar 39.95
4	19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.90	23 V 50.94	24 Cr 52.01	25 Mn 54.9	26 Fe 55.85	27 Co 58.71	28 Ni 58.71	29 Cu 63.54	30 Zn 65.37	31 Ga 69.7	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.91	36 Kr 83.80	
5	37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 91.22	42 Mo 95.94	43 Tc 98.9	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3	
6	55 Cs 132.9	56 Ba 137.3	71 Lu 174.9	72 Hf 178.5	73 Ta 180.9	74 W 183.8	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 196.9	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 208.9	84 Po 210	85 At 210	86 Rn 222	
7	87 Fr 223	88 Ra 226.0	103 Lr 257	104 Unq	105 Unp	106 Unh	107 Uns	108 Uno	109 Une										

Lanthanides		57 La 138.9	58 Ce 140.1	59 Pr 140.9	60 Nd 144.2	61 Pm 146.9	62 Sm 150.9	63 Eu 151.3	64 Gd 157.3	65 Tb 158.9	66 Dy 162.5	67 Ho 164.9	68 Er 167.3	69 Tm 168.9	70 Yb 173.0		
Actinides		89 Ac 227.0	90 Th 232.0	91 Pa 231.0	92 U 238.0	93 Np 237.1	94 Pu 239.1	95 Am 241.1	96 Cm 247.1	97 Bk 249.1	98 Cf 251.1	99 Es 254.1	100 Fm 257.1	101 Md 258.1	102 No 255		

Numbers below the symbol indicates the atomic masses; and the numbers above the symbol indicates the atomic numbers.

SOURCE: International Union of Pure and Applied Chemistry, I mills, ed., Quantities, Units, and symbols in Physical Chemistry, Blackwell Scientific publications, Boston, 1988, pp 86-98.

Useful Relations				General Data	
$(RT)_{298.15K} = 2.4789 \text{ kJ/mol}$				speed of light	$c = 2.997\ 925 \times 10^8 \text{ ms}^{-1}$
$(RT/F)_{298.15K} = 0.025\ 693 \text{ V}$				charge of proton	$e = 1.602\ 19 \times 10^{-19} \text{ C}$
T/K: 100.15 298.15 500.15 1000.15				Faraday constant	$F = 9.648\ 46 \times 10^4 \text{ C mol}^{-1}$
T/Cm ⁻¹ : 69.61 207.22 347.62 695.13				Boltzmann constant	$k = 1.380\ 66 \times 10^{-23} \text{ J K}^{-1}$
1mmHg=133.222 N m ⁻²				Gas constant	$R = 8.314\ 41 \text{ J K}^{-1} \text{ mol}^{-1}$
hc/k=1.438 78x10 ⁻² m K					$8.205\ 75 \times 10^{-2} \text{ dm}^3 \text{ atm K}^{-1} \text{ mol}^{-1}$
SI-units:					
$1 \text{ L} = 1000 \text{ ml} = 1000 \text{ cm}^3 = 1 \text{ dm}^3$				Planck constant	$h = 6.626\ 18 \times 10^{-34} \text{ Js}$
$=1.01325 \times 10^5 \text{ Nm}^{-2}$	$=4.184 \text{ J}$	$=1.602\ 189 \times 10^{-19} \text{ J}$	$=0.124 \times 10^{-3} \text{ eV}$		$\hbar = \frac{h}{2\pi} = 1.054\ 59 \times 10^{-34} \text{ Js}$
$=760 \text{ torr}$		$=96.485 \text{ kJ/mol}$	$=1.9864 \times 10^{-23} \text{ J}$		
$=1 \text{ bar}$		$=8065.5 \text{ cm}^{-1}$		Avogadro constant	$N_A = 6.022\ 14 \times 10^{23} \text{ mol}^{-1}$
				Atomis mass unit	$u = 1.660\ 54 \times 10^{-27} \text{ kg}$
				Electron mass	$m_e = 9.109\ 39 \times 10^{-31} \text{ kg}$
				Proton mass	$m_p = 1.672\ 62 \times 10^{-27} \text{ kg}$
				Neutron mass	$m_n = 1.674\ 93 \times 10^{-27} \text{ kg}$
				Vacuum permittivity	$\epsilon_0 = \mu_0^{-1} c^{-2} = 8.854\ 188 \times 10^{-12} \text{ J}^{-1} \text{ C}^2 \text{ m}^{-1}$
dipole moment: 1 Debye = $3.335\ 64 \times 10^{-30} \text{ C m}$				Vacuum permeability	$\mu_0 = 4\pi \times 10^{-7} \text{ Js}^2 \text{ C}^{-2} \text{ m}^{-1}$
force: $1N = 1 \text{ J m}^{-1} = 1 \text{ kg ms}^{-2} = 10^5 \text{ dyne}$	pressure: $1Pa = 1 \text{ N m}^{-2} = 1 \text{ J m}^{-3}$			Bohr magneton	$\mu_B = \frac{e\hbar}{2m_e} = 9.274\ 02 \times 10^{-24} \text{ JT}^{-1}$
$1 \text{ J} = 1 \text{ N m}$					
power: $1 \text{ W} = 1 \text{ J s}^{-1}$	potential: $1 \text{ V} = 1 \text{ J C}^{-1}$			Nuclear magneton	$\mu_N = \frac{e\hbar}{2m_p} = 5.05079 \times 10^{-27} \text{ JT}^{-1}$
magnetic flux: $1 \text{ T} = 1 \text{ V sm}^{-2} = 1 \text{ J C sm}^{-2}$	current: $1 \text{ A} = 1 \text{ C s}^{-1}$			Gravitational constant	$G = 6.67259 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
Prefixes:				Gravitational acceleration	$g = 9.80665 \text{ ms}^{-2}$
p n m m c d k M G					
pico nano micro milli centi deci kilo mega giga				Bohr radius	$a_0 = 5.291\ 77 \times 10^{-11} \text{ m}$
10^{-12}	10^{-9}	10^{-6}	10^{-3}		
10^{-2}	10^{-1}	10^3	10^6		

Standard molar enthalpies of formation at 298.15 K

Temperature dependence of heat capacities, $C_{p,m} = a+bT+cT^{-2}$

	M_r	$\Delta H_f^\theta / \text{kJ/mol}$		M_r	$\Delta H_f^\theta / \text{kJ/mol}$		$a/\text{J K}^{-1}\text{mol}^{-1}$	$b/10^{-3}\text{J K}^{-2}\text{mol}^{-1}$	$c/10^5\text{J K}\text{mol}^{-1}$
H ₂ O(g)	18.015	-241.8	O ₃ (g)	47.998	+142.7	Gases (298-2000K)			
H ₂ O(l)	18.015	-285.8	NO(g)	30.006	+90.2	He, Ne, Ar, Kr, Xe	20.78	0	0
H ₂ O ₂ (l)	34.015	-187.8	NO ₂ (g)	46.006	+33.2	H ₂	27.28	3.26	0.50
NH ₃ (g)	17.031	-46.1	N ₂ O ₄ (g)	92.012	+9.2	O ₂	29.96	4.18	-1.67
N ₂ H ₄ (l)	32.045	+50.6	SO ₂ (g)	64.063	-296.8	N ₂	28.58	3.77	-0.50
N ₃ H(l)	43.028	+264.1	H ₂ S(g)	34.080	-20.6	Cl ₂	37.03	0.67	-2.85
N ₃ H(g)	43.028	+294.1	SF ₆ (g)	146.054	-1209	CO ₂	44.23	8.79	-8.62
HNO ₃ (l)	63.013	-174.1	HF(g)	20.006	-271.1	H ₂ O	30.54	10.29	0
NH ₃ OH(s)	33.030	-114.2	HCl(g)	36.461	-92.3	NH ₃	29.75	25.10	-1.55
NH ₄ Cl(s)	53.492	-314.4	HCl(aq)	36.461	-167.2	CH ₄	23.64	47.86	-1.92
HgCl ₂ (s)	271.50	-224.3	HBr(g)	80.917	+36.4				
H ₂ SO ₄ (l)	98.078	-814.0	Hl(g)	127.912	+26.5				
H ₂ SO ₄ (aq)	98.078	-909.3	CO ₂ (g)	44.010	-393.5				
NaCl(s)	58.443	-411.0	CO(g)	28.011	-110.5				
NaOH(s)	39.997	-426.7	Al ₂ O ₃ (α ,s)	101.945	-1675.7				
KCl(s)	74.555	-435.9	SiO ₂ (s)	60.085	-910.9				
KBr(s)	119.011	-392.2	FeS(s)	87.91	-100.0				
KI(s)	166.006	-327.6	FeS ₂ (s)	119.975	-178.2				
Diatomics(g)	---	0	AgCl(s)	143.323	-127.1				
Enthalpies of fusion and evaporation $\Delta H_m^\theta / \text{kJ/mol}$ at the transition temperature									
T _f /K	Fusion ^a	T _b /K	Evaporation ^b				M_r	$\Delta H_f^\theta / \text{kJ/mol}$	$\Delta H_c^\theta / \text{kJ/mol}$
He	3.5	0.021	4.22	0.084			CH ₄ (g)	16.043	-74.81
Ar	83.81	1.188	87.29	6.506			C ₂ H ₂ (g)	26.038	+226.8
H ₂	13.96	0.117	20.38	0.9163			C ₂ H ₄ (g)	28.054	+52.30
N ₂	63.15	0.719	77.35	5.586			C ₂ H ₆ (g)	30.070	-84.64
O ₂	54.36	0.444	90.18	6.820			C ₃ H ₆ cyclopropane(g)	42.081	53.35
Cl ₂	172.12	6.406	239.05	20.410			C ₃ H ₆ (propene(g))	42.081	20.5
Br ₂	265.90	10.573	332.35	29.45			C ₄ H ₁₀ n-butane (g)	58.124	-126.11
I ₂	386.75	15.52	458.39	41.80			C ₅ H ₁₂ n-pentane(g)	72.151	-146.4
Hg	234.29	2.292	629.73	59.296			C ₆ H ₁₂ cyclohexane (l)	84.163	-156.2
Ag	1234	11.30	2436	250.63			C ₆ H ₁₄ n-hexane (l)	86.178	-198.7
Na	370.95	2.601	1156	98.01			C ₆ H ₆ benzene (l)	78.115	+48.99
CO ₂	217.0	8.33	194.64	25.23	↓		C ₈ H ₁₈ n-octane (l)	114.233	-249.8
H ₂ O	273.15	6.008	373.15	40.656	(44.016 at 298.15 K)		C ₁₀ H ₈ naphthalene (l)	128.175	+78.53
NH ₃	195.40	5.652	239.73	23.351			CH ₃ OH (l)	32.042	-239.0
H ₂ S	187.61	2.377	212.80	18.673			CH ₃ CHO (g)	44.054	-166.0
CH ₄	90.68	0.941	111.66	8.18			CH ₃ CH ₂ OH (l)	46.070	-277.0
C ₂ H ₆	89.85	2.86	184.55	14.7			CH ₃ COOH (l)	60.053	-484.2
C ₆ H ₆	278.65	10.59	353.25	30.8			CH ₃ COOC ₂ H ₅ (l)	88.107	-486.6
CH ₃ OH	175.25	3.159	337.22	35.27	(37.99 at 298.15K)		C ₆ H ₅ OH (s)	94.114	-165.0
							C ₆ H ₅ NH ₂ (l)	93.129	-31.1
							NH ₂ CO.NH, urea(s)	60.056	-333.0
							CH ₂ (NH ₂)CO ₂ H, glycine (s)	75.068	-537.2
							C ₆ H ₁₂ O ₆ , α -D-glucose (s)	180.159	-1274
							C ₆ H ₂₂ O ₆ , β -D-glucose (s)	180.159	-1268
							C ₁₂ H ₂₂ O ₁₁ , sucrose (s)	342.303	-2222
							CH ₃ CH(OH)COOH	90.079	-694.0
							lactic acid (s)		1344

↓ Sublimation: ^avarious pressures: ^bat 1atm

Source: American Institute of Physics handbook, McGraw-Hill

Standard molar Gibbs free energy and molar entropy of formation at 298.15 K

	M _r	ΔG _f ^θ /kJ/mol	S ^θ /J K ⁻¹ mol ⁻¹		M _r	ΔG _f ^θ /kJ/mol	S ^θ /J K ⁻¹ mol ⁻¹
H ₂ O(g)	18.015	-228.57	188.83	O ₃ (g)	47.998	163.2	238.93
H ₂ O(l)	18.015	-120.35	109.6	NO(g)	30.006	86.55	210.76
H ₂ O ₂ (l)	34.015	-120.35	109.6	NO ₂ (g)	46.006	51.31	240.06
NH ₃ (g)	17.031	-16.45	192.45	N ₂ O ₄ (g)	92.012	97.89	304.29
N ₂ H ₄ (l)	32.045	149.43	121.21	SO ₂ (g)	64.063	-300.19	248.22
N ₃ H(l)	43.028	327.3	140.6	H ₂ S(g)	34.080	-33.56	205.79
N ₃ H(g)	43.028	328.1	238.97	SF ₆ (g)	146.054	-1105.3	291.82
HNO ₃ (l)	63.013	-80.71	155.60	HF(g)	20.006	-273.2	173.78
NH ₄ OH(s)	33.030			HCl(g)	36.461	-95.30	186.91
NH ₄ Cl(s)	53.492	-202.87	94.6	HCl(aq)	36.461	-131.23	56.5
HgCl ₂ (s)	271.50	-178.6	146.0	HBr(g)	80.917	-53.45	198.70
H ₂ SO ₄ (l)	98.078	-690.00	156.90	HI(g)	127.912	1.70	206.59
H ₂ SO ₄ (aq)	98.078	-744.53	20.1	CO ₂ (g)	44.010	-394.36	213.74
NaCl(s)	58.443	-384.14	72.13	CO(g)	28.011	-137.17	197.67
NaOH(s)	39.997	-379.49	64.46	Al ₂ O ₃ (s)	101.945	-1582.3	50.92
KCl(s)	74.555	-409.14	82.59	SiO ₂	60.09	-856.64	41.84
KBr(s)	119.011	-380.66	95.90	FeS(s)	87.91	-100.4	60.29
KI(s)	166.006	-324.89	106.32	FeS ₂ (s)	119.975	-166.9	52.93
				AgCl(s)	143.323	-109.79	96.2
He(g)	4.003	0	126.15	Hg(g)	200.59	31.82	174.96
Ar(g)	39.95	0	154.84	Hg(l)	200.59	0	76.02
H ₂ (g)	2.016	0	130.684	Ag(g)	107.87	245.65	173.00
N ₂ (g)	28.013	0	191.61	Ag(s)	107.87	0	42.55
O ₂ (g)	31.999	0	205.138	Na(g)	370.95	76.76	153.71
O ₃ (g)	47.998	163.2	238.93	Na(s)	22.99	0	51.21
Cl ₂ (g)	70.91	0	223.07				
Br ₂ (g)	159.82	3.110	245.46				
Br ₂ (l)	159.82	0	152.23				
I ₂ (g)	253.81	19.33	260.69				
I ₂ (s)	253.81	0	116.135				

	M _r	ΔG _f ^θ /kJ/mol	S ^θ /J K ⁻¹ mol ⁻¹
organic compounds			
CH ₄ (g) methane	16.043	-50.72	186.26
C ₂ H ₂ (g) ethyne	26.038	209.20	200.94
C ₂ H ₄ (g) ethene	28.05	68.15	219.56
C ₂ H ₆ (g) ethane	30.070	-32.82	229.60
C ₃ H ₆ cyclopropane(g)	42.081	104.45	237.55
C ₃ H ₆ propene(g)	42.081	62.78	267.05
C ₄ H ₁₀ n-butane (g)	58.124	-17.03	310.23
C ₅ H ₁₂ n-pentane(g)	72.151	-8.20	348.40
C ₆ H ₁₂ cyclohexane (l)	84.163	26.8	
C ₆ H ₁₄ n-hexane (l)	86.178		204.3
C ₆ H ₆ benzene (l)	78.115	124.3	173.3
C ₆ H ₆ benzene (g)	78.115	129.72	269.31
C ₈ H ₁₈ n-octane (l)	114.233	6.4	361.1
C ₁₀ H ₈ naphthalene (l)	128.175		
CH ₃ OH (g)	32.042	-161.96	239.81
CH ₃ OH (l)	32.042	-166.27	126.8
CH ₃ CHO (g)	44.054	-128.86	250.3
CH ₃ CH ₂ OH (l)	46.07	-174.78	160.7
CH ₃ COOH (l)	60.053	-389.9	159.8
CH ₃ COOC ₂ H ₅ (l)	88.107	-332.7	259.4
C ₆ H ₅ OH (s)	94.114	-50.9	146.0
C ₆ H ₅ NH ₂ (l)	93.129		
CH ₂ (NH ₂)CO ₂ H, glycine (s)	75.068	-373.4	103.5
C ₆ H ₁₂ O ₆ , b-D-glucose (s)	180.159	-910	212
C ₁₂ H ₂₂ O ₁₁ , sucrose (s)	342.303	-1543	360.2
CH ₃ CH(OH)COOH	90.079		
lactic acid (s)			

Source: American Institute of Physics handbook, McGraw-Hill.

Heat capacities at 25°C

	C _{v,m} JK ⁻¹ mol ⁻¹	C _{p,m} JK ⁻¹ mol ⁻¹
He, Ne, Ar, Kr, Xe	12.47	20.78
H ₂	20.50	28.81
O ₂	21.01	29.33
N ₂	20.83	29.14
CO ₂	28.83	37.14
NH ₃	27.17	35.48
CH ₄	27.43	35.74
N ₂ O ₄		77.28
NO ₂		37.20

F.P Depression, B.P. Elevation

Solvent	F.P °C	K _f °C kg mol ⁻¹	B.P (°C, 101kNm ⁻²)	K _b °C kg mol ⁻¹
Water	0	1.86	100.0	0.52
Benzene	5.51	5.10	80.1	2.60
Acetic Acid	16.6	3.90	118.1	3.10
Cyclohexane	6.5	20.2	81.4	2.79
Camphor	177.7	40.0	205	-
Nitrobenzene	5.7	6.9	210.9	5.24
Ethanol	-177		78.5	1.22
Chloroform	-64		61.3	3.63

Third Law entropies at 25°C, Sm^θ/J K⁻¹ mol⁻¹

Solids	Liquids	Gases
Ag	42.68	H ₂
C(gr)	5.77	N ₂
C(d)	2.44	O ₂
Cu	33.4	Cl ₂
Zn	41.6	
I ₂	116.7	CO ₂
S(Rh)	31.9	HCl
		H ₂ S
AgCl	96.2	NH ₃
AgBr	104.6	CH ₄
CuSO ₄ ·5H ₂ O	305.4	C ₂ H ₆
HgCl ₂	144	CH ₃ CHO
Sucrose	360.2	
	C ₂ H ₅ OH	
	CH ₃ OH	
	C ₆ H ₆	
	CH ₃ COOH	
	C ₆ H ₁₂	