

UNIVERSITY OF SWAZILAND
BACHELOR OF SCIENCE

SUPPLEMENTARY/RESIT EXAMINATION 2018

TITLE OF PAPER : INTRODUCTION TO THERMODYNAMICS /
(COURSE CODE) (CHE241)

INTRODUCTORY PHYSICAL CHEMISTRY
(C202)

TIME : 3 HOURS

INSTRUCTIONS : THERE ARE SIX QUESTIONS

: ANSWER ANY FOUR QUESTIONS

: BEGIN THE ANSWER TO EACH QUESTION ON
A SEPARATE SHEET OF PAPER

: DATA SHEETS ARE PROVIDED WITH THIS
EXAMINATION PAPER

DO NOT OPEN THIS PAPER UNTIL THE INVIGILATOR INSTRUCTS YOU TO DO SO.

Question 1 [25 Marks]

- a) Gases behave as ideal and non-ideal in defined P,V and T zones. With the aid of **any two** of the following: [15]
- Lennard-Jones potential plot;
 - Compressibility plots and
 - Isotherm plots,

Compare and contrast real and ideal gases. Your account should make mention of interactions, equations and the kinetic theory of gases to help clarify your discussion.

- b) A mixture of butane (C_4H_{10}) and propene (C_3H_6) occupied 35.5 L at 1.000 bar and 405 K. This mixture **reacted completely** with 220.6 g of O_2 to produce CO_2 and H_2O .
- What was the composition of the original mixture? Assume ideal gas behaviour.
MW (O_2)=32 g/mol [5]
 - Calculate the partial pressure, mole fraction of each gas and the total pressure of the final mixture. [5]

Question 2 [25 Marks]

A real gas equation of state for a gas is given by:

$$P = \frac{RT}{V_m - nb} - \frac{a}{V_m^2} \quad (1)$$

- a) Based on van der Waals assumptions discuss the bases and significance of the main terms in equation (1) in terms of gas behaviour. [10]
- b) Derive expressions for $V_{m,c}$, T_c and P_c . [6]
- c) Find an expression for the Boyle's temperature, T_B . [5]
- d) Estimate the temperature at which oxygen behaves as an ideal gas, T_B given the constants:
 $a=6.493 \text{ L}^2\text{atmmol}^{-2}$, $b=5.622 \times 10^{-2} \text{ Lmol}^{-1}$ [2]
- e) Estimate the radii of real gas molecules using equation (1) for real gases given a critical molar volume of $250 \text{ cm}^3\text{mol}^{-1}$ [2]

Question 3 [25 Marks]

- a) Using examples and/or diagrams compare and contrast **Any Two (2)** of the following terms
- Reversible and irreversible expansion [5]
 - Path and state functions [5]
 - Work and heat [5]
- b) the work done during the isothermal reversible expansion of a gas that satisfies the virial equation of state
Evaluate:

$$\frac{PV_m}{RT} = 1 + \frac{B}{V_m} + \frac{C}{V^2} + \dots; B = -21.7 \text{ cm}^3 \text{ mol}^{-1} \text{ & } C = 1200 \text{ cm}^6 \text{ mol}^{-2}$$

- (i). Derive an expression for work for a real gas that satisfies the virial equation in a reversible isothermal expansion [6]
- (ii). Calculate work for 1.0 mol Ar at 273 K obeying the virial gas equation [4]
- (iii). Calculate work for 1.0 mol Ar at 273 K obeying the perfect gas equation. [5]

Let expansion be from 500 cm³ to 1000 cm³ in each case.

QUESTION 4 [25 marks]


Adiabatic expansion of an ideal gas is quite different from isothermal expansion.

- a) Explain what is meant by adiabatic expansion, draw an adiabat and an isotherm on a P versus V graph and compare them. [8]
- b) Derive the expression for the change in temperature of an adiabatic expansion of an ideal gas against constant external pressure from V_i to V_f. [8]
- c) A sample of argon at 1.0 atm pressure and 25°C expands reversibly and adiabatically from 0.50 L to 1.00 L. calculate: [9]
 - i) final temperature
 - ii) work done
 - iii) change in internal energy.

QUESTION 5 [25 MARKS]

- a) Write short notes on any two of the following
 - i) Enthalpy change [5]
 - ii) Internal energy change [5]
 - iii) Hess's Law [5]
 - b) To Calibrate a calorimeter a 0.120 g naphthalene, C₁₀H₈(s), was burned at constant volume and it caused the temperature of the calorimeter to rise by 3.05 K. Then 0.10 g of an unknown compound was burned in the same calorimeter, causing a temperature rise of 2.05 K.
 - i) Calculate the heat capacity of the calorimeter [3]
 - ii) Is the unknown compound phenol, C₆H₅OH(s) or ethanol, CH₃CH₂OH(l) whose enthalpies of combustion are Δ_CH^θ=-3054 kJmol⁻¹ and -1368 kJmol⁻¹ respectively. [4]
 - c) Calculate the standard enthalpies of formation of:
 - i) KClO₃(s) from the enthalpy of formation of KCl [4]
 - ii) NOCl(g) from the enthalpy of formation of NO [4]
- Given the attached table and the following information:





$$\Delta_f H^\theta = +76.5 \text{ kJ/mol}$$

Question 6 [25 Marks]

- a) Briefly discuss the statistical view of entropy [5]
- b) 1.00 mol of perfect gas at 27°C is expanded isothermally from an initial pressure of 3.00 atm to a final pressure of 1.00 atm. Calculate q , w , ΔS_{sys} , ΔS_{surr} and ΔS_{tot} if the expansion is done:
- (1) reversibly. [5]
 - (2) against a constant external pressure of 1.00 atm. [5]
 - (3) adiabatically against a constant pressure of 1.00 atm. [5]
- c) If 50g water at 80°C is poured into 100g water at 10°C in an insulated vessel given that $C_{\text{p,m}}=75.5 \text{ JK}^{-1}\text{mol}^{-1}$; Calculate:
- i) final temperature of the mixture [3]
 - ii) the entropy change [2]
-

Useful Relations				General Data	
$(RT)_{298.15K} = 2.4789 \text{ kJ/mol}$				speed of light	$c = 2.997\ 925 \times 10^8 \text{ ms}^{-1}$
$(RT/F)_{298.15K} = 0.025\ 693 \text{ V}$				charge of proton	$e = 1.602\ 19 \times 10^{-19} \text{ C}$
T/K: 100.15 298.15 500.15 1000.15				Faraday constant	$F = Le = 9.648\ 46 \times 10^4 \text{ C mol}^{-1}$
T/Cm ⁻¹ : 69.61 207.22 347.62 695.13				Boltzmann constant	$k = 1.380\ 66 \times 10^{-23} \text{ J K}^{-1}$
1mmHg=133.222 N m ⁻²				Gas constant	$R = Lk = 8.314\ 41 \text{ J K}^{-1} \text{ mol}^{-1}$
hc/k=1.438 78x10 ⁻² m K					$8.205\ 75 \times 10^{-2} \text{ dm}^3 \text{ atm K}^{-1} \text{ mol}^{-1}$
1atm	1 cal	1 eV	1cm ⁻¹		
= $1.01325 \times 10^5 \text{ Nm}^{-2}$	=4.184 J	= $1.602\ 189 \times 10^{-19} \text{ J}$	= $0.124 \times 10^{-3} \text{ eV}$	Planck constant	$h = 6.626\ 18 \times 10^{-34} \text{ Js}$
=760torr		=96.485 kJ/mol	= $1.9864 \times 10^{-23} \text{ J}$		$\hbar = \frac{h}{2\pi} = 1.054\ 59 \times 10^{-34} \text{ Js}$
=1 bar		= 8065.5 cm ⁻¹			
SI-units:					
$1 \text{ L} = 1000 \text{ ml} = 1000 \text{ cm}^3 = 1 \text{ dm}^3$				Avogadro constant	$L \text{ or } N_{av} = 6.022\ 14 \times 10^{23} \text{ mol}^{-1}$
$1 \text{ dm} = 0.1 \text{ m}$				Atomis mass unit	$u = 1.660\ 54 \times 10^{-27} \text{ kg}$
$1 \text{ cal (thermochemical)} = 4.184 \text{ J}$				Electron mass	$m_e = 9.109\ 39 \times 10^{-31} \text{ kg}$
dipole moment: $1 \text{ Debye} = 3.335\ 64 \times 10^{-30} \text{ C m}$				Proton mass	$m_p = 1.672\ 62 \times 10^{-27} \text{ kg}$
force: $1N = 1 \text{ J m}^{-1} = 1 \text{ kgms}^{-2} = 10^5 \text{ dyne}$ pressure: $1Pa = 1 \text{ Nm}^{-2} = 1 \text{ J m}^{-3}$				Neutron mass	$m_n = 1.674\ 93 \times 10^{-27} \text{ kg}$
$1J = 1 \text{ Nm}$				Vacuum permittivity	$\epsilon_0 = \mu_0^{-1} c^2 = 8.854\ 188 \times 10^{-12} \text{ J}^{-1} \text{ C}^2 \text{ m}^{-1}$
power: $1W = 1 \text{ J s}^{-1}$			potential: $1V = 1 \text{ J C}^{-1}$	Vacuum permeability	$\mu_0 = 4\pi \times 10^{-7} \text{ Js}^2 \text{ C}^{-2} \text{ m}^{-1}$
				Bohr magneton	$\mu_B = \frac{e\hbar}{2m_e} = 9.274\ 02 \times 10^{-24} \text{ JT}^{-1}$
magnetic flux: $1T = 1 \text{ Vs m}^{-2} = 1 \text{ JCsm}^{-2}$			current: $1A = 1 \text{ Cs}^{-1}$	Nuclear magneton	$\mu_N = \frac{e\hbar}{2m_p} = 5.05079 \times 10^{-27} \text{ JT}^{-1}$
Prefixes:				Gravitational constant	$G = 6.67259 \times 10^{-11} \text{ Nm}^2 \text{ kg}^{-2}$
p n m m c d k M G				Gravitational acceleration	$g = 9.80665 \text{ ms}^{-2}$
pico nano micro milli centi deci kilo mega giga				Bohr radius	$a_0 = 5.291\ 77 \times 10^{-11} \text{ m}$
10^{-12}	10^{-9}	10^{-6}	10^{-3}		
			10^{-2}		
			10^{-1}		
			10^3		
			10^6		
			10^9		

Heat capacities at 25°C

	C _{v,m}	C _{p,m}
	JK ⁻¹ mol ⁻¹	JK ⁻¹ mol ⁻¹
He, Ne, Ar, Kr, Xe	12.47	20.78
H ₂	20.50	28.81
O ₂	21.01	29.33
N ₂	20.83	29.14
CO ₂	28.83	37.14
NH ₃	27.17	35.48
CH ₄	27.43	35.74
N ₂ O ₄		77.28
NO ₂		37.20

F.P Depression, B.P. Elevation

Solvent	F.P °C	K _f °C kg mol ⁻¹	B.P (°C, 101kNm ⁻²)	K _b °C kg mol ⁻¹
Water	0	1.86	100.0	0.52
Benzene	5.51	5.10	80.1	2.60
Acetic Acid	16.6	3.90	118.1	3.10
Cyclohexane	6.5	20.2	81.4	2.79
Camphor	177.7	40.0	205	-
Nitrobenzene	5.7	6.9	210.9	5.24
Ethanol	-177		78.5	1.22
Chloroform	-64		61.3	3.63

Third Law entropies at 25°C, Sm^θ/J K⁻¹ mol⁻¹

Solids	Liquids	Gases
Ag	42.68	H ₂
C(gr)	5.77	N ₂
C(d)	2.44	O ₂
Cu	33.4	Cl ₂
Zn	41.6	
I ₂	116.7	CO ₂
S(Rh)	31.9	HCl
AgCl	96.2	H ₂ S
AgBr	104.6	NH ₃
CuSO ₄ ·5H ₂ O	305.4	CH ₄
HgCl ₂	144	C ₂ H ₆
Sucrose	360.2	CH ₃ CHO
	C ₆ H ₁₂	229.4
		265.7

Standard molar enthalpies of formation at 298.15 K

Temperature dependence of heat capacities, $C_{p,m} = a+bT+cT^{-2}$

	M_r	$\Delta H_f^\theta / \text{kJ/mol}$		M_r	$\Delta H_f^\theta / \text{kJ/mol}$		$a/\text{J K}^{-1}\text{mol}^{-1}$	$b/10^{-3}\text{J K}^{-2}\text{mol}^{-1}$	$c/10^5\text{J K}\text{mol}^{-1}$
H ₂ O(g)	18.015	-241.8	O ₃ (g)	47.998	+142.7	Gases (298-2000K)			
H ₂ O(l)	18.015	-285.8	NO(g)	30.006	+90.2	He, Ne, Ar, Kr, Xe	20.78	0	0
H ₂ O ₂ (l)	34.015	-187.8	NO ₂ (g)	46.006	+33.2	H ₂	27.28	3.26	0.50
NH ₃ (g)	17.031	-46.1	N ₂ O ₄ (g)	92.012	+9.2	O ₂	29.96	4.18	-1.67
N ₂ H ₄ (l)	32.045	+50.6	SO ₂ (g)	64.063	-296.8	N ₂	28.58	3.77	-0.50
N ₂ H(l)	43.028	+264.1	H ₂ S(g)	34.080	-20.6	Cl ₂	37.03	0.67	-2.85
N ₂ H(g)	43.028	+294.1	SF ₆ (g)	146.054	-1209	CO ₂	44.23	8.79	-8.62
HNO ₃ (l)	63.013	-174.1	HF(g)	20.006	-271.1	H ₂ O	30.54	10.29	0
NH ₄ OH(s)	33.030	-114.2	HCl(g)	36.461	-92.3	NH ₃	29.75	25.10	-1.55
NH ₄ Cl(s)	53.492	-314.4	HCl(aq)	36.461	-167.2	CH ₄	23.64	47.86	-1.92
HgCl ₂ (s)	271.50	-224.3	HBr(g)	80.917	+36.4				
H ₂ SO ₄ (l)	98.078	-814.0	Hf(g)	127.912	+26.5				
H ₂ SO ₄ (aq)	98.078	-909.3	CO ₂ (g)	44.010	-393.5				
NaCl(s)	58.443	-411.0	CO(g)	28.011	-110.5				
NaOH(s)	39.997	-426.7	Al ₂ O ₃ (α , s)	101.945	-1675.7				
KCl(s)	74.555	-435.9	SiO ₂ (s)	60.085	-910.9				
KBr(s)	119.011	-392.2	FeS(s)	87.91	-100.0				
KI(s)	166.006	-327.6	FeS ₂ (s)	119.975	-178.2				
Diatomics(g)	—	0	AgCl(s)	143.323	-127.1				
						M_r	$\Delta H_f^\theta / \text{kJ/mol}$	$\Delta H_c^\theta / \text{kJ/mol}$	
						CH ₄ (g)	16.043	-74.81	
						C ₂ H ₂ (g)	26.038	+226.8	1300
						C ₂ H ₄ (g)	28.054	+52.30	1411
						C ₂ H ₆ (g)	30.070	-84.64	1560
						C ₃ H ₆ cyclopropane(g)	42.081	53.35	2091
						C ₃ H ₆ propene(g)	42.081	20.5	2058
						C ₄ H ₁₀ n-butane (g)	58.124	-126.11	2877
						C ₅ H ₁₂ n-pentane(g)	72.151	-146.4	3536
						C ₆ H ₁₂ cyclohexane (l)	84.163	-156.2	3920
						C ₆ H ₁₄ n-hexane (l)	86.178	-198.7	4163
						C ₆ H ₆ benzene (l)	78.115	+48.99	3268
						C ₈ H ₁₈ n-octane (l)	114.233	-249.8	5471
						C ₁₀ H ₈ naphthalene (l)	128.175	+78.53	5157
						CH ₃ OH (l)	32.042	-239.0	726.1
						CH ₃ CHO (g)	44.054	-166.0	1193
						CH ₃ CH ₂ OH (l)	46.070	-277.0	1368
						CH ₃ COOH (l)	60.053	-484.2	874.5
						CH ₃ COOC ₂ H ₅ (l)	88.107	-486.6	2231
						C ₆ H ₅ OH (s)	94.114	-165.0	3054
						C ₆ H ₅ NH ₂ (l)	93.129	-31.1	3393
						NH ₂ CO.NH, urea(s)	60.056	-333.0	632.2
						CH ₂ (NH ₂)CO ₂ H, glycine (s)	75.068	-537.2	964.4
						C ₆ H ₁₂ O ₆ , α -D-glucose (s)	180.159	-1274	2802
						C ₆ H ₂₂ O ₆ , β -D-glucose (s)	180.159	-1268	2808
						C ₁₂ H ₂₂ O ₁₁ , sucrose (s)	342.303	-2222	5645
						CH ₃ CH(OH)COOH	90.079	-694.0	1344
						lactic acid (s)			

Sublimation: ^avarious pressures: ^bat 1atm

Source: American Institute of Physics handbook, McGraw-Hill

Standard molar Gibbs free energy and molar entropy of formation at 298.15 K

	M _r	$\Delta G_f^\theta/\text{kJ/mol}$	$S^\theta/\text{J K}^{-1}\text{ mol}^{-1}$		M _r	$\Delta G_f^\theta/\text{kJ/mol}$	$S^\theta/\text{J K}^{-1}\text{ mol}^{-1}$
H ₂ O(g)	18.015	-228.57	188.83	O ₃ (g)	47.998	163.2	238.93
H ₂ O(l)	18.015	-120.35	109.6	NO(g)	30.006	86.55	210.76
H ₂ O ₂ (l)	34.015	-120.35	109.6	NO ₂ (g)	46.006	51.31	240.06
NH ₃ (g)	17.031	-16.45	192.45	N ₂ O ₄ (g)	92.012	97.89	304.29
N ₂ H ₄ (l)	32.045	149.43	121.21	SO ₂ (g)	64.063	-300.19	248.22
N ₂ H(l)	43.028	327.3	140.6	H ₂ S(g)	34.080	-33.56	205.79
N ₂ H(g)	43.028	328.1	238.97	SF ₆ (g)	146.054	-1105.3	291.82
HNO ₃ (l)	63.013	-80.71	155.60	HF(g)	20.006	-273.2	173.78
NH ₄ OH(s)	33.030			HCl(g)	36.461	-95.30	186.91
NH ₄ Cl(s)	53.492	-202.87	94.6	HCl(aq)	36.461	-131.23	56.5
HgCl ₂ (s)	271.50	-178.6	146.0	HBr(g)	80.917	-53.45	198.70
H ₂ SO ₄ (l)	98.078	-690.00	156.90	HI(g)	127.912	1.70	206.59
H ₂ SO ₄ (aq)	98.078	-744.53	20.1	CO ₂ (g)	44.010	-394.36	213.74
NaCl(s)	58.443	-384.14	72.13	CO(g)	28.011	-137.17	197.67
NaOH(s)	39.997	-379.49	64.46	Al ₂ O ₃ (s)	101.945	-1582.3	50.92
KCl(s)	74.555	-409.14	82.59	SiO ₂	60.09	-856.64	41.84
KBr(s)	119.011	-380.66	95.90	FeS(s)	87.91	-100.4	60.29
KI(s)	166.006	-324.89	106.32	FeS ₂ (s)	119.975	-166.9	52.93
				AgCl(s)	143.323	-109.79	96.2
He(g)	4.003	0	126.15	Hg(g)	200.59	31.82	174.96
Ar(g)	39.95	0	154.84	Hg(l)	200.59	0	76.02
H ₂ (g)	2.016	0	130.684	Ag(g)	107.87	245.65	173.00
N ₂ (g)	28.013	0	191.61	Ag(s)	107.87	0	42.55
O ₂ (g)	31.999	0	205.138	Na(g)	370.95	76.76	153.71
O ₃ (g)	47.998	163.2	238.93	Na(s)	22.99	0	51.21
Cl ₂ (g)	70.91	0	223.07				
Br ₂ (g)	159.82	3.110	245.46				
Br ₂ (l)	159.82	0	152.23				
I ₂ (g)	253.81	19.33	260.69				
I ₂ (s)	253.81	0	116.135				

	M _r	$\Delta G_f^\theta/\text{kJ/mol}$	$S^\theta/\text{J K}^{-1}\text{ mol}^{-1}$
organic compounds			
CH ₄ (g) methane	16.043	-50.72	186.26
C ₂ H ₂ (g) ethyne	26.038	209.20	200.94
C ₂ H ₄ (g) ethene	28.05	68.15	219.56
C ₂ H ₆ (g) ethane	30.070	-32.82	229.60
C ₃ H ₆ cyclopropane(g)	42.081	104.45	237.55
C ₃ H ₆ propene(g)	42.081	62.78	267.05
C ₄ H ₁₀ n-butane (g)	58.124	-17.03	310.23
C ₅ H ₁₂ n-pentane(g)	72.151	-8.20	348.40
C ₆ H ₁₂ cyclohexane (l)	84.163	26.8	
C ₆ H ₁₄ n-hexane (l)	86.178		204.3
C ₆ H ₆ benzene (l)	78.115	124.3	173.3
C ₆ H ₆ benzene (g)	78.115	129.72	269.31
C ₈ H ₁₈ n-octane (l)	114.233	6.4	361.1
C ₁₀ H ₈ naphthalene (l)	128.175		
CH ₃ OH (g)	32.042	-161.96	239.81
CH ₃ OH (l)	32.042	-166.27	126.8
CH ₃ CHO (g)	44.054	-128.86	250.3
CH ₃ CH ₂ OH (l)	46.07	-174.78	160.7
CH ₃ COOH (l)	60.053	-389.9	159.8
CH ₃ COOC ₂ H ₅ (l)	88.107	-332.7	259.4
C ₆ H ₅ OH (s)	94.114	-50.9	146.0
C ₆ H ₅ NH ₂ (l)	93.129		
CH ₂ (NH ₂)CO ₂ H, glycine (s)	75.068	-373.4	103.5
C ₆ H ₁₂ O ₆ , b-D-glucose (s)	180.159	-910	212
C ₁₂ H ₂₂ O ₁₁ , sucrose (s)	342.303	-1543	360.2
CH ₃ CH(OH)COOH	90.079		
lactic acid (s)			

Source: American Institute of Physics handbook, McGraw-Hill.

THE PERIODIC TABLE OF ELEMENTS

Group	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
	IA	IIA	IIIB	IVB	VIB	VIB	VIIB	VIIIB			IB	IIB	IIIA	IVA	VA	VIA	VIIA	VIIIA
Period 1	1 H 1.008																	2 He 4.003
2	3 Li 6.94	4 Be 9.01																10 Ne 20.18
3	11 Na 22.99	12 Mg 24.31																18 Ar 39.95
4	19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.90	23 V 50.94	24 Cr 52.01	25 Mn 54.9	26 Fe 55.85	27 Co 58.71	28 Ni 58.71	29 Cu 63.54	30 Zn 65.37	31 Ga 69.7	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.91	36 Kr 83.80
5	37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 91.22	42 Mo 95.94	43 Tc 98.9	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3
6	55 Cs 132.9	56 Ba 137.3	71 Lu 174.9	72 Hf 178.5	73 Ta 180.9	74 W 183.8	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 196.9	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 208.9	84 Po 210	85 At 210	86 Rn 222
7	87 Fr 223	88 Ra 226.0	103 Lr 257	104 Unq	105 Unp	106 Unh	107 Uns	108 Uno	109 Une									

Lanthanides	57 La 138.9	58 Ce 140.1	59 Pr 140.9	60 Nd 144.2	61 Pm 146.9	62 Sm 150.9	63 Eu 151.3	64 Gd 157.3	65 Tb 158.9	66 Dy 162.5	67 Ho 164.9	68 Er 167.3	69 Tm 168.9	70 Yb 173.0
Actinides	89 Ac 227.0	90 Th 232.0	91 Pa 231.0	92 U 238.0	93 Np 237.1	94 Pu 239.1	95 Am 241.1	96 Cm 247.1	97 Bk 249.1	98 Cf 251.1	99 Es 254.1	100 Fm 257.1	101 Md 258.1	102 No 255

Numbers below the symbol indicates the atomic masses; and the numbers above the symbol indicates the atomic numbers.

SOURCE: International Union of Pure and Applied Chemistry, I mills, ed., Quantities, Units, and symbols in Physical Chemistry, Blackwell Scientific publications, Boston, 1988, pp 86-98.