

**UNIVERSITY OF ESWATINI**

**MAIN EXAMINATION 2018/2019**

**TITLE OF PAPER:** **INTRODUCTION TO QUANTUM MECHANICS**

**COURSE NUMBER:** **CHE343**

**TIME:** **THREE (3) HOURS**

**INSTRUCTIONS:**

**Answer all questions**

**NB:** Each question should start on a new page.

A data sheet and a periodic table are attached

A non-programmable electronic calculator may be used

**DO NOT OPEN THIS PAPER UNTIL PERMISSION TO DO SO HAS BEEN GRANTED  
BY THE CHIEF INVIGILATOR.**

### **QUESTION 1 [25 MARKS]**

- a) A particle is moving in one dimension between  $x = 1\text{cm}$  and  $x = 2\text{cm}$ . the potential is such that the particle cannot be found outside these limits and the wave function in between is  $\psi = B/x$
- Find the normalization constant [5]
  - Calculate the average value of  $x$  [5]
- b) Which of the following functions below are eigenfunctions of the operator  $d^2/dx^2$ ?  
For each eigenfunction give the eigenvalue.
- $\exp(ikx)$
  - $6\cos(4x)$
  - $\exp(-ax^2)$
  - $k^2$  [5]
- c) the x component of angular momentum of an orbiting electron is  $L_x = zp_x - xp_z$ .
- Find the quantum mechanical operator  $\hat{L}_x$  [2]
  - Evaluate the following commutator:  $[\hat{Y}, \hat{P}_y]$  [3]
- d) For a particle in a box whose ends are at  $x = 0$  and  $x = 0.4000\text{nm}$ , calculate the probability that the particle's x coordinate is between  $0.16000$  and  $0.16001\text{nm}$  if  $n=1$  [5]
- $$\left[ \psi = \left( \frac{2}{L} \right)^{\frac{1}{2}} \sin\left( \frac{n\pi x}{L} \right) \right]$$

### **QUESTION 2 [25 MARKS]**

Given that a particle in a one dimensional box has the wavefunctionn

$$\psi = \left( \frac{2}{L} \right)^{\frac{1}{2}} \sin\left( \frac{n\pi x}{L} \right), \quad n=1,2,3\dots \text{ for } 0 < x < L$$

- Derive the expression for the energy of the particle [7]
- If the particle was now confined in a 3 dimensional cubic box, what would be the energy expression? [4]

c) If now the cubical box has dimensions  $L_x = L_y = L_z/3$ , what would be the energy when

- i).  $n_x = 1, n_y = 2, n_z = 2$
- ii).  $n_x = 1, n_y = 1, n_z = 4$

What values of  $n_x, n_y$  and  $n_z$  would make the two energy levels degenerate? [7]

d) .

- i). Distinguish between the terms “**state**” and “**energy level**” of a system.
- ii). For a particle in a cubic box of edge  $L$ , how many states have the energies in the range 0 to  $20\hbar^2/(8mL^2)$ ?
- iii). How many energy levels lie in this range? [7]

### **QUESTION 3 (25 MARKS)**

a) How did the study of heat capacity of metals consolidate the Plank's hypothesis that energy is quantized? [8]

b) .

- i). Write down the expression for a one dimensional harmonic oscillator, defining all terms. [3]
- ii). Assuming the vibrations of  $^{14}\text{N}_2$  molecule are equivalent to those of a harmonic oscillator with a force constant  $k = 2293.8 \text{ N/m}$ , what is the zero point energy of the vibrations of this molecule. (the mass of the nitrogen molecule is  $14.004\text{u}$ ) [7]

c) The rotation of  $\text{H}^{127}\text{I}$  molecule can be pictured as the orbital motion of an H atom at a distance 0.160 nm from a stationary **iodine** atom. Suppose the molecule rotates only in one plane.

- i). Calculate the energy needed to excite the molecule into rotation [4]
- ii). What is the minimum non-zero angular momentum of the molecule? [3]

**Total Marks**

**/75/**

## Useful Integrals

- 
1.  $\int x^2 e^{-x^2} dx = \frac{\sqrt{\pi}}{2}$
  2.  $\int x^3 e^{-x^2} dx = 0$
  3.  $\int_0^\infty x^n e^{-ax} dx = \frac{n!}{a^{n+1}}$
  4.  $\int \sin \theta d\theta = -\cos \theta + \text{constant}$
  5.  $d\tau = r^2 \sin \theta dr d\theta d\phi$
  6.  $\int x^n dx = \frac{1}{a^{n+1}} \quad n \neq -1$
  7.  $\int_0^{2\pi} \cos^2 \theta \sin \theta d\theta = \frac{2}{3}$
-

## General data and fundamental constants

Quantity	Symbol	Value
Speed of light	c	$2.997\ 924\ 58 \times 10^8 \text{ m s}^{-1}$
Elementary charge	e	$1.602\ 177 \times 10^{-19} \text{ C}$
Faraday constant	F = $N_A e$	$9.6485 \times 10^4 \text{ C mol}^{-1}$
Boltzmann constant	k	$1.380\ 66 \times 10^{-23} \text{ J K}^{-1}$
Gas constant	R = $N_A k$	$8.314\ 51 \text{ J K}^{-1} \text{ mol}^{-1}$
		$8.205\ 78 \times 10^{-2} \text{ dm}^3 \text{ atm K}^{-1} \text{ mol}^{-1}$
Planck constant	h	$6.2364 \times 10^{-34} \text{ J s}$
	$\hbar = h/2\pi$	$6.626\ 08 \times 10^{-34} \text{ J s}$
Avogadro constant	$N_A$	$1.054\ 57 \times 10^{23} \text{ J s}$
Atomic mass unit	u	$6.022\ 14 \times 10^{23} \text{ mol}^{-1}$
Mass		$1.660\ 54 \times 10^{-27} \text{ Kg}$
electron	$m_e$	$9.109\ 39 \times 10^{-31} \text{ Kg}$
proton	$m_p$	$1.672\ 62 \times 10^{-27} \text{ Kg}$
neutron	$m_n$	$1.674\ 93 \times 10^{-27} \text{ Kg}$
Vacuum permittivity	$\epsilon_0 = 1/c^2 \mu_0$	$8.854\ 19 \times 10^{-12} \text{ J}^{-1} \text{ C}^2 \text{ m}^{-1}$
	$4\pi\epsilon_0$	$1.112\ 65 \times 10^{-10} \text{ J}^{-1} \text{ C}^2 \text{ m}^{-1}$
Vacuum permeability	$\mu_0$	$4\pi \times 10^{-7} \text{ J s}^{-2} \text{ C}^{-2} \text{ m}^{-1}$
		$4\pi \times 10^{-7} \text{ T}^2 \text{ J}^{-1} \text{ m}^3$
Magneton		
Bohr	$\mu_B = e\hbar/2m_e$	$9.274\ 02 \times 10^{-24} \text{ J T}^{-1}$
nuclear	$\mu_N = e\hbar/2m_p$	$5.050\ 79 \times 10^{-27} \text{ J T}^{-1}$
g value	$g_e$	2.002 32
Bohr radius	$a_0 = 4\pi\epsilon_0\hbar/m_e e^2$	$5.291\ 77 \times 10^{-11} \text{ m}$
Fine-structure constant	$\alpha = \mu_0 e^2 c / 2h$	$7.297\ 35 \times 10^{-3}$
Rydberg constant	$R_\infty = m_e e^4 / 8\hbar^3 c \epsilon_0^2$	$1.097\ 37 \times 10^7 \text{ m}^{-1}$
Standard acceleration of free fall	g	$9.806\ 65 \text{ m s}^{-2}$
Gravitational constant	G	$6.672\ 59 \times 10^{-11} \text{ N m}^2 \text{ Kg}^{-2}$

## Conversion factors

$$\begin{aligned}
 1 \text{ cal} &= 4.184 \text{ joules (J)} \\
 1 \text{ eV} &= 1.602 2 \times 10^{-19} \text{ J}
 \end{aligned}
 \quad
 \begin{aligned}
 1 \text{ erg} &= 1 \times 10^{-7} \text{ J} \\
 1 \text{ eV/molecule} &= 96 485 \text{ kJ mol}^{-1}
 \end{aligned}$$

Prefixes	f	p	n	$\mu$	m	c	d	k	M	G
	femto	pico	nano	micro	milli	centi	deci	kilo	mega	giga
	$10^{-15}$	$10^{-12}$	$10^{-9}$	$10^{-6}$	$10^{-3}$	$10^{-2}$	$10^{-1}$	$10^3$	$10^6$	$10^9$

# PERIODIC TABLE OF ELEMENTS

## GROUPS

PERIODS	GROUPS																									
	1A 1.008	2A 11	3A 12	4A 13	5A 14	6A 15	7A 16	VIB 17	VIIIB 18	VIIIB 19	VIIIB 20	VIIIB 21	VIIIB 22	VIIIB 23	VIIIB 24	VIIIB 25	VIIIB 26	VIIIB 27	VIIIB 28	VIIIB 29	VIIIB 30	VIIIB 31	VIIIB 32	VIIIB 33	VIIIB 34	VIIIB 35
1	6.941 Li 3	9.012 Be 4																								
2	22.990 Na 11	24.305 Mg 12																								
3	39.098 K 19	40.078 Ca 20	44.956 Sc 21	47.88 Ti 22	50.942 V 23	51.996 Cr 24	54.938 Mn 25	55.847 Fe 26	58.933 Co 27	58.69 Ni 28	63.546 Cu 29	65.39 Zn 30	72.61 Ga 31	74.922 Ge 32	78.96 As 33	79.904 Se 34	83.80 Br 35	Kr 36								
4	85.468 Rb 37	87.62 Sr 38	88.906 Y 39	91.224 Zr 40	92.906 Nb 41	95.94 Mo 42	98.907 Tc 43	101.07 Ru 44	102.91 Rh 45	106.42 Pd 46	107.87 Ag 47	112.41 Cd 48	114.82 In 49	118.71 Sn 50	121.75 Sb 51	127.60 Te 52	126.90 I 53	Xe 54								
5	137.33 Cs 55	137.91 Ba 56	138.91 *La 57	178.49 Hf 72	180.95 Ta 73	183.85 W 74	186.21 Re 75	190.2 Os 76	192.22 Ir 77	195.08 Pt 78	196.97 Au 79	198.98 Hg 80	207.2 Tl 81	208.98 Pb 82	(209)	(210)	(222)	Rn 85	86							
6	223 Fr 87	226.03 Ra 88	(227) **Ac 89	(261) Rf 104	(262) Ha 105	(263) Unh 106	(262) Unh 107	(265) Uns 108	(266) Uno 109	(267) Une 110																
7																										

\* Lanthanide Series

\*\* Actinide Series

140.12 Ce 58	140.91 Pr 59	144.24 Nd 60	(145) Pm 61	150.36 Sm 62	151.96 Eu 63	157.25 Gd 64	158.93 Tb 65	162.50 Dy 66	164.93 Ho 67	167.26 Er 68	168.93 Tm 69	173.04 Yb 70	174.97 Lu 71
232.04 Th 90	231.04 Pa 91	238.03 U 92	237.05 Np 93	(244) Pu 94	(243) Am 95	(247) Cm 96	(247) Bk 97	(251) Cf 98	(252) Ds 99	(257) Fm 100	(258) Md 101	(259) No 102	(260) Lr 103

( ) indicates the mass number of the isotope with the longest half-life.