

UNIVERSITY OF SWAZILAND
SUPPLEMENTARY EXAMINATION PAPER 2007

TITLE OF PAPER: QUANTITATIVE METHODS IN DEMOGRAPHY

COURSE CODE : DEM 206

TIME ALLOWED : TWO (2) HOURS

**INSTRUCTIONS : THIS PAPER HAS FIVE QUESTIONS. ANSWER
ANY THREE (3) QUESTIONS.**

**PLEASE DO NOT OPEN THIS PAPER UNTIL PERMISSION HAS BEEN
GRANTED BY THE INVIGILATOR**

QUESTION 1 (8+6+6 marks)

- a. The probability that a man will be alive in 25 years is $\frac{3}{5}$, and the probability that his wife will be alive in 25 years is $\frac{2}{3}$. Find the probability that in 25 years:
- both will be alive;
 - only the man will be alive;
 - only the wife will be alive; and
 - at least one will be alive.
- b. Out of 500 families with 4 children each, what percentage would be expected to have:
- 2 boys and 2 girls;
 - at least one boy; and
 - at most 2 girls?
- Assume equal probabilities for boys and girls.
- c. A recent study of robberies for a certain geographic region showed an average of one robbery per 20,000 people. In a city of 80,000 people, find the probability of the following:
- At least one robbery?
 - At most two robberies?

QUESTION 2 (5+5+5+5 marks)

Random samples of size 3 are drawn from the finite population which consists of the numbers 5, 6, 7, 8, 9, and 10.

- Compute the population mean and standard deviation.
- List all the possible random samples of size 3 that can be drawn from this finite population and calculate their means.
- Construct the sampling distribution of the mean for random samples of size 3 from this given population.
- Calculate the mean and standard error of the sampling distribution.

QUESTION 3 (14+6 marks)

- a. A statistician claims that the average age of people who purchase lottery tickets is 70. A sample of 30 is selected, and their ages are recorded. At $\alpha = 0.05$, is there enough evidence to reject the statistician's claim?

49	80	24	61	79	68
63	72	46	65	76	91
90	56	70	71	71	67
52	82	74	39	49	69
22	56	70	74	62	45

- b. Construct an interval estimate of the average age of people who purchase lottery tickets. Use a confidence level of 95 %.

QUESTION 4 (12+8 marks)

Write short notes on the following:

- Estimation of trend in time series analysis.
- Approaches to measuring probability.

QUESTION 5 (8+4+8 marks)


- a. A study is done to see whether there is a relationship between a mother's age and the number of children she has. The data are shown below.

Number of children	2	1	3	1	2	4	3	5
Mother's age	18	22	29	20	27	32	33	36

- Predict the number of children of a mother whose age is 34.
- How important is age in explaining the differences in the number of children the women bear?

b. A study on crime suggested that, at least 40% of all arsonists were under 21 years of age. Checking local crime statistics, a researcher found that 30 out of 80 arson suspects were under 21 years. At $\alpha = 0.01$, should the crime statistics be rejected?

Table 4. Normal curve areas
Standard normal probability in right-hand tail
(for negative values of z areas are found by symmetry)



z	Second decimal place of z								
	.00	.01	.02	.03	.04	.05	.06	.07	.08
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681
0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286
0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897
0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520
0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156
0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810
0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483
0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177
0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894
0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635
1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401
1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190
1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003
1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0837
1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0722	.0708	.0694
1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571
1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465
1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375
1.8	.0359	.0352	.0344	.0336	.0329	.0322	.0314	.0307	.0301
1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239
2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188
2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146
2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113
2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087
2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066
2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049
2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037
2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027
2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0019
2.9	.0019	.0018	.0017	.0017	.0016	.0016	.0015	.0015	.0014
3.0	.00135								
3.5	.000 233								
4.0	.000 031 7								
4.5	.000 003 40								
5.0	.000 000 287								

From R. R. Walpole, *Introduction to Statistics* (New York: Macmillan, 1968).

Table 5. ~~Quantiles~~ Percentage points of the χ^2 -distribution

$\chi^2_{0.100}$	$\chi^2_{0.050}$	$\chi^2_{0.025}$	$\chi^2_{0.010}$	$\chi^2_{0.005}$	d.f.
2.70554	3.84146	5.02389	6.63490	7.87944	1
4.60517	5.99147	7.37776	9.21034	10.5966	2
6.25139	7.81473	9.34840	11.3449	12.8381	3
7.77944	9.48773	11.1433	13.2767	14.8602	4
9.23635	11.0705	12.8325	15.0863	16.7496	5
10.6446	12.5916	14.4494	16.8119	18.5476	6
12.0170	14.0671	16.0128	18.4753	20.2777	7
13.3616	15.5073	17.5346	20.0902	21.9550	8
14.6837	16.9190	19.0228	21.6660	23.5893	9
15.9871	18.3070	20.4831	23.2093	25.1882	10
17.2750	19.6751	21.9200	24.7250	26.7569	11
18.5494	21.0261	23.3367	26.2170	28.2995	12
19.8119	22.3621	24.7356	27.6883	29.8194	13
21.0642	23.6848	26.1190	29.1413	31.3193	14
22.3072	24.9958	27.4884	30.5779	32.8013	15
23.5418	26.2962	28.8454	31.9999	34.2672	16
24.7690	27.5871	30.1910	33.4087	35.7185	17
25.9894	28.8693	31.5264	34.8053	37.1564	18
27.2036	30.1435	32.8523	36.1908	38.5822	19
28.4120	31.4104	34.1696	37.5662	39.9968	20
29.6151	32.6705	35.4789	38.9321	41.4010	21
30.8133	33.9244	36.7807	40.2894	42.7956	22
32.0069	35.1725	38.0757	41.6384	44.1813	23
33.1963	36.4151	39.3641	42.9798	45.5585	24
34.3816	37.6525	40.6465	44.3141	46.9278	25
35.5631	38.8852	41.9232	45.6417	48.2899	26
36.7412	40.1133	43.1944	46.9630	49.6449	27
37.9159	41.3372	44.4607	48.2782	50.9933	28
39.0875	42.5569	45.7222	49.5879	52.3356	29
40.2560	43.7729	46.9792	50.8922	53.6720	30
51.8050	55.7585	59.3417	63.6907	66.7659	40
63.1671	67.5048	71.4202	76.1539	79.4900	50
74.3970	79.0819	83.2976	88.3794	91.9517	60
85.5271	90.5312	95.0231	100.425	104.215	70
96.5782	101.879	106.629	112.329	116.321	80
107.565	113.145	118.136	124.116	128.299	90
118.498	124.342	129.561	135.807	140.169	100

From "Tables of the Percentage Points of the χ^2 -Distribution," *Biometrika*, Vol. 32 (1941), pp. 188-189, by Catherine M. Thompson. Reproduced by permission of Professor R. S. Pearson.