

UNIVERSITY OF SWAZILAND

FINAL EXAMINATION PAPER 2007

TITLE OF PAPER : MULTIVARIATE ANALYSIS

COURSE CODE : ST410

TIME ALLOWED : 2 (TWO) HOURS

**REQUIRMENTS : STATISTICAL TABLES
AND CALCULATOR**

**INSTRUCTIONS : ANSWER ANY 4 (FOUR) QUESTIONS.
ALL QUESTIONS CARRY EQUAL MARKS.**

**THIS PAPER IS NOT TO BE OPENED UNTIL PERMISSION HAS BEEN
GRANTED BY THE INVIGILATOR**

QUESTION ONE.

[3 + 7 + 7 + 6 + 2 marks]

Suppose we have three variables in each of the 3 groups with sample sizes $n_A=3$, $n_B=4$ and $n_C=5$. Consider the followings:

$$\bar{x} = \begin{bmatrix} 4 & 13 & 5 \\ 6 & 10 & 6 \\ 3 & 12 & 7 \end{bmatrix}, \quad \bar{X} = \begin{pmatrix} 4.25 \\ 11.58 \\ 6.17 \end{pmatrix}, \quad S^2 = \begin{pmatrix} 5.7 \\ 8.6 \\ 1.6 \end{pmatrix}, \quad C^{-1} = \begin{bmatrix} 2.170 & -0.258 & -2.435 \\ -0.258 & 0.156 & 0.148 \\ -2.435 & 0.148 & 3.643 \end{bmatrix}$$
$$W = \begin{bmatrix} 42.0 & 39.0 & 10.0 \\ 39.0 & 78.0 & 11.0 \\ 10.0 & 11.0 & 10.0 \end{bmatrix}, \quad \& \quad T = \begin{bmatrix} 62.25 & 24.25 & 8.5 \\ 24.25 & 94.92 & 8.83 \\ 8.5 & 8.83 & 17.67 \end{bmatrix}$$

where $\bar{x}$ is the matrix of means, the first row represents the means of the three variables in group A, etc.; $\bar{X}$ is the vector of means of the three variables; S^2 is the vector of variances of the three variables; C^{-1} is the inverse of the pooled covariance matrix of group A and C; W is the within sum of square matrix and T is the total sum of square matrix.

- 1.1 Perform a univariate t tests for variables 2 and 3 at 5% level of significance.
- 1.2 Perform Hotellings' T^2 test considering groups B and C.
- 1.3 Compute Wilk's Λ statistic and ϕ . Use χ^2 approximation to test the equality of population mean vectors. Specify the null and alternative hypotheses.
- 1.4 Complete the ANOVA tables, with one extra column for the conclusion, of the one-way analysis of variance procedure to test the equality of variable means for each of those three variables.
- 1.5 Comment on the results found in part (1.3) and (1.4).

QUESTION TWO.

[10 + 8 + 2 + 2 + 3 marks]

- 2.1 Discuss the importance of Principal Component Analysis and state its important properties.
- 2.2 Explain the procedure for a principal component analysis.
- 2.3 Consider the following table:

Eigenvalue	Eigenvectors						
	X_1	X_2	X_3	X_4	X_5	X_6	X_7
3.111	0.512	0.375	-0.246	-0.315	-0.222	-0.382	-0.131
1.709	-0.024	0.000	0.432	0.109	-0.242	-0.408	-0.553
1.095	-0.278	0.516	-0.503	-0.292	0.071	0.064	-0.096
0.663	0.016	0.113	0.058	0.023	0.783	0.169	-0.489
0.305	0.025	-0.345	0.231	-0.854	-0.064	0.269	-0.133
0.108	-0.045	0.203	-0.028	0.208	-0.503	0.674	-0.399
0.009	0.166	-0.212	-0.238	0.065	0.014	-0.165	-0.463

- a. How many variables were there in the data set? How many components will you get?
- b. How many components will you choose? Explain why.
- c. List those selected components and interpret those in terms of original variables, X_i 's.

QUESTION THREE.

[6 + 2 + 3 + 3 + 3 + 8 marks]

The following tables are part of the complete output running SPSS for a set of multivariate variables; not necessarily from the same set of variables. Tables 1-6 are obtained running Factor Analysis and Tables 7-9 are obtained running Discriminant Function Analysis:

Table 1:

Correlation Matrix							
	X1	X2	X3	X4	X5	X6	X7
Correlatio X1	1.000	.036	-.671	-.400	-.538	-.645	-.764
X2	.036	1.000	.445	.405	-.026	-.495	-.221
X3	-.671	.445	1.000	.385	.494	.080	.200
X4	-.400	.405	.385	1.000	.060	.199	.185
X5	-.538	-.026	.494	.060	1.000	.271	.210
X6	-.645	-.495	.080	.199	.271	1.000	.424
X7	-.764	-.221	.200	.185	.210	.424	1.000

Table 2:

Component	Eigenvalues
1	2.964
2	1.950
3	0.871
4	0.643
5	0.359
6	0.211
7	0.001

Table 3:

Component Matrix ^a						
	Component					
	1	2	3	4	5	6
X1	-.986	5.070E-02	-2.47E-02	.113	.102	-2.99E-02
X2	-1.24E-02	.936	7.822E-02	-.109	3.541E-04	.325
X3	.683	.566	-.258	-3.93E-02	-.333	-.186
X4	.483	.481	.590	.327	.241	-.153
X5	.621	4.503E-02	-.677	.186	.344	4.014E-02
X6	.641	-.562	.179	.402	-.186	.212
X7	.705	-.319	.246	-.560	.166	2.041E-02

Extraction Method: Principal Component Analysis.
a. 6 components extracted.

Table 4:

Component Matrix ^a			
	Component		
	1	2	3
X1	-.986	5.070E-02	-2.47E-02
X2	-1.24E-02	.936	7.822E-02
X3	.683	.566	-.258
X4	.483	.481	.590
X5	.621	4.503E-02	-.677
X6	.641	-.562	.179
X7	.705	-.319	.246

Extraction Method: Principal Component Analysis.
a. 3 components extracted.

Table 5:

Rotated Component Matrix ^a			
	Component		
	1	2	3
X1	-.764	-.550	-.301
X2	-.515	.195	.761
X3	8.552E-02	.756	.523
X4	.295	-2.72E-02	.851
X5	.185	.892	-.127
X6	.858	8.626E-02	-.124
X7	.790	.141	.120

Extraction Method: Principal Component Analysis.
Rotation Method: Varimax with Kaiser Normalization.
a. Rotation converged in 6 iterations.

Table 6:

	Component		
	1	2	3
X1	-.270	-.170	-.103
X2	-.263	.078	.433
X3	-.102	.422	.169
X4	.192	-.291	.584
X5	-.113	.658	-.290
X6	.400	-.097	-.063
X7	.365	-.099	.083

Extraction Method: Principal Component Analysis.
Rotation Method: Varimax with Kaiser Normalization.

Table 7:

Wilks' Lambda				
Test of Function(s)	Wilks' Lambda	Chi-square	df	Sig.
1 through 2	.147	39.244	12	.000
2	.656	8.655	5	.124

Table 8:

Standardized Canonical Discriminant Function Coefficients

	Function	
	1	2
X1	.268	1.193
X2	.202	-1.065
X3	.412	.938
X4	.573	.478
X5	.129	.097
X6	-1.040	-.249

Table 9:

Canonical Discriminant Function Coefficients

	Function	
	1	2
X1	.019	.083
X2	.305	-1.603
X3	.059	.135
X4	1.512	1.263
X5	.076	.057
X6	-.268	-.064
(Constant)	.217	-3.732

Unstandardized coefficients

- 3.1 Examine Tables 1 & 2 and explain the suitability of principal component analysis. How many principal components will you obtain from the original data set? How many principal components will you choose? Explain.
- 3.2 How many factors will you choose if you wish to use factor analysis method? Explain your answer.
- 3.3 How many factors will you get in your factor model from Table 3? List the last equation of your model and compute its communality.
- 3.4 Suppose the same data were analyzed using with a restriction on the number of factors and the results are given in Table 4. What was restricted number of factors? List the first two equations of your model and compute their communalities.
- 3.5 List all equations needed to compute factor scores.
- 3.6 Write all the discriminant functions and test whether each of those is significant at 5% level of significance.

QUESTION FOUR.

[4 + 5 + 4 + 4 + 8 marks]

- 4.1 Explain why the Discriminant Function Analysis is different from the Principal Component Analysis and the Factor Analysis.
- 4.2 Discuss the procedures to obtain canonical discriminant functions.
- 4.3 The following table shows the eigenvalues and corresponding eigenvectors of $W^{-1}B$:

Component	Eigenvalue	Eigenvectors			
		X_1	X_2	X_3	X_4
1	0.437	-0.0107	0.0040	0.0119	-0.0068
2	0.035	0.0031	0.0168	-0.0046	-0.0022
3	0.015	-0.0068	0.0010	0.0000	0.0247
4	0.002	0.0126	-0.0001	0.0112	0.0054

- a. How many groups and variables were considered in this problem?
- b. List all the canonical discriminant functions.
- c. Assuming that the i^{th} sample size, $n_i = 32$ for all $i = 1, 2, 3, 4, 5$; test whether each of these functions varies significantly from group to group.

QUESTION FIVE.

[15 + 3 + 7 marks]

- 5.1 What is Factor Model? Explain how to obtain principal component factors.
- 5.2 Why do we need to use factor rotation to obtain final factors?
- 5.3 Write the unrotated factor model along with the respective communalities using the following table which shows the eigenvalues and corresponding eigenvectors of C^{-1} :

Component	Eigenvalue	Eigenvectors				
		X_1	X_2	X_3	X_4	X_5
1	2.616	0.452	0.462	0.451	0.471	0.398
2	1.532	-0.051	0.300	0.325	0.185	-0.377
3	0.386	0.691	0.341	-0.455	-0.411	-0.179
4	0.302	-0.420	0.548	-0.606	0.388	0.069
5	0.165	0.374	-0.530	-0.343	0.652	-0.192

Table 5
Percentage points of the t distributions




$t_{.100}$	$t_{.050}$	$t_{.025}$	$t_{.010}$	$t_{.005}$	df
3.078	6.314	12.706	31.821	63.657	1
1.886	2.920	4.303	6.965	9.925	2
1.638	2.353	3.182	4.541	5.841	3
1.533	2.132	2.776	3.747	4.604	4
1.476	2.015	2.571	3.365	4.032	5
1.440	1.943	2.447	3.143	3.707	6
1.415	1.895	2.365	2.998	3.499	7
1.397	1.860	2.306	2.896	3.355	8
1.383	1.833	2.262	2.821	3.250	9
1.372	1.812	2.228	2.764	3.169	10
1.363	1.796	2.201	2.718	3.106	11
1.356	1.782	2.179	2.681	3.055	12
1.350	1.771	2.160	2.650	3.012	13
1.345	1.761	2.145	2.624	2.977	14
1.341	1.753	2.131	2.602	2.947	15
1.337	1.746	2.120	2.583	2.921	16
1.333	1.740	2.110	2.567	2.898	17
1.330	1.734	2.101	2.552	2.878	18
1.328	1.729	2.093	2.539	2.861	19
1.325	1.725	2.086	2.528	2.845	20
1.323	1.721	2.080	2.518	2.831	21
1.321	1.717	2.074	2.508	2.819	22
1.319	1.714	2.069	2.500	2.807	23
1.318	1.711	2.064	2.492	2.797	24
1.316	1.708	2.060	2.485	2.787	25
1.315	1.706	2.056	2.479	2.779	26
1.314	1.703	2.052	2.473	2.771	27
1.313	1.701	2.048	2.467	2.763	28
1.311	1.699	2.045	2.462	2.756	29
1.282	1.645	1.960	2.326	2.576	inf.

From "Table of Percentage Points of the t-Distribution,"
Computed by Maxine Merrington, *Biometrika*, Vol. 32 (1941), p.
300. Reproduced by permission of Professor E. S. Pearson.

TABLE A.3 Percentiles of the χ^2 Distribution

Entry is $\chi^2(A; \nu)$ where $P\{\chi^2(\nu) \leq \chi^2(A; \nu)\} = A$




ν	.005	.010	.025	.050	.100	.900	.950	.975	.990	.995
1	0.00393	0.0157	0.00982	0.02393	0.0158	2.71	3.84	5.02	6.63	7.88
2	0.0100	0.0201	0.0506	0.103	0.211	4.61	5.99	7.38	9.21	10.60
3	0.072	0.115	0.216	0.352	0.584	6.25	7.81	9.35	11.34	12.84
4	0.207	0.297	0.484	0.711	1.064	7.78	9.49	11.14	13.28	14.86
5	0.412	0.554	0.831	1.145	1.61	9.24	11.07	12.83	15.09	16.75
6	0.676	0.872	1.24	1.64	2.20	10.64	12.59	14.45	16.81	18.55
7	0.989	1.24	1.69	2.17	2.83	12.02	14.07	16.01	18.48	20.28
8	1.34	1.65	2.18	2.73	3.49	13.36	15.51	17.53	20.09	21.96
9	1.73	2.09	2.70	3.33	4.17	14.68	16.92	19.02	21.67	23.59
10	2.16	2.56	3.25	3.94	4.87	15.99	18.31	20.48	23.21	25.19
11	2.60	3.05	3.82	4.57	5.58	17.28	19.68	21.92	24.73	26.76
12	3.07	3.57	4.40	5.23	6.30	18.55	21.03	23.34	26.22	28.30
13	3.57	4.11	5.01	5.89	7.04	19.81	22.36	24.74	27.69	29.82
14	4.07	4.66	5.63	6.57	7.79	21.06	23.68	26.12	29.14	31.32
15	4.60	5.23	6.26	7.26	8.55	22.31	25.00	27.49	30.58	32.80
16	5.14	5.81	6.91	7.96	9.31	23.54	26.30	28.85	32.00	34.27
17	5.70	6.41	7.56	8.67	10.09	24.77	27.59	30.19	33.41	35.72
18	6.26	7.01	8.23	9.39	10.86	25.99	28.87	31.53	34.81	37.16
19	6.84	7.63	8.91	10.12	11.65	27.20	30.14	32.85	36.19	38.58
20	7.43	8.26	9.59	10.85	12.44	28.41	31.41	34.17	37.57	40.00
21	8.03	8.90	10.28	11.59	13.24	29.62	32.67	35.48	38.93	41.40
22	8.64	9.54	10.98	12.34	14.04	30.81	33.92	36.78	40.29	42.80
23	9.26	10.20	11.69	13.09	14.85	32.01	35.17	38.08	41.64	44.18
24	9.89	10.86	12.40	13.85	15.66	33.20	36.42	39.36	42.98	45.56
25	10.52	11.52	13.12	14.61	16.47	34.38	37.65	40.65	44.31	46.93
26	11.16	12.20	13.84	15.38	17.29	35.56	38.89	41.92	45.64	48.29
27	11.81	12.88	14.57	16.15	18.11	36.74	40.11	43.19	46.96	49.64
28	12.46	13.56	15.31	16.93	18.94	37.92	41.34	44.46	48.28	50.99
29	13.12	14.26	16.05	17.71	19.77	39.09	42.56	45.72	49.59	52.34
30	13.79	14.95	16.79	18.49	20.60	40.26	43.77	46.98	50.89	53.67
40	20.71	22.16	24.43	26.51	29.05	51.81	55.76	59.34	63.69	66.77
50	27.99	29.71	32.36	34.76	37.69	63.17	67.50	71.42	76.15	79.49
60	35.53	37.48	40.48	43.19	46.46	74.40	79.08	83.30	88.38	91.95
70	43.28	45.44	48.76	51.74	55.33	85.53	90.53	95.02	100.4	104.2
80	51.17	53.54	57.15	60.39	64.28	96.58	101.9	106.6	112.3	116.3
90	59.20	61.75	65.65	69.13	73.29	107.6	113.1	118.1	124.1	128.3
100	67.33	70.06	74.22	77.93	82.36	118.5	124.3	129.6	135.8	140.2

Source: Reprinted, with permission, from C. M. Thompson, "Table of Percentage Points of the Chi-Square Distribution," *Biometrika* 32 (1941), pp. 188-89.

TABLE A.4 Percentiles of the F Distribution

Entry is $F(A; \nu_1, \nu_2)$ where $P\{F(\nu_1, \nu_2) \leq F(A; \nu_1, \nu_2)\} = A$



$F(A; \nu_1, \nu_2) = \frac{F(1-A; \nu_2, \nu_1)}{1 - A}$



TABLE A.4 (continued) Percentiles of the F Distribution

Den. df	A	Numerator df								
		1	2	3	4	5	6	7	8	9
1	.50	1.00	1.50	1.71	1.82	1.89	1.94	1.98	2.00	2.03
	.90	39.9	49.5	53.6	55.8	57.2	58.2	58.9	59.4	59.9
	.95	161	200	216	225	230	234	237	239	241
	.975	648	800	864	900	922	937	948	957	963
	.99	4,032	5,000	5,403	5,625	5,764	5,859	5,928	5,981	6,022
	.995	16,211	20,000	21,615	22,500	23,056	23,437	23,715	23,925	24,091
2	.50	0.667	1.00	1.13	1.21	1.25	1.28	1.30	1.32	1.33
	.90	8.53	9.00	9.16	9.24	9.29	9.33	9.35	9.37	9.38
	.95	18.5	19.0	19.2	19.2	19.3	19.3	19.4	19.4	19.4
	.975	38.5	39.0	39.2	39.2	39.3	39.3	39.4	39.4	39.4
	.99	98.5	99.0	99.2	99.2	99.3	99.3	99.4	99.4	99.4
	.995	199	199	199	199	199	199	199	199	199
3	.50	0.585	0.881	1.00	1.06	1.10	1.13	1.15	1.16	1.17
	.90	5.54	5.46	5.39	5.34	5.31	5.28	5.27	5.25	5.24
	.95	10.1	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81
	.975	17.4	16.0	15.4	15.1	14.9	14.7	14.6	14.5	14.5
	.99	34.1	30.8	29.5	28.7	28.2	27.9	27.7	27.5	27.3
	.995	55.6	49.8	47.5	46.2	45.4	44.8	44.4	44.1	43.9
4	.50	0.549	0.828	0.941	1.00	1.04	1.06	1.08	1.09	1.10
	.90	4.54	4.32	4.19	4.11	4.05	4.01	3.98	3.95	3.94
	.95	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00
	.975	12.2	10.6	9.98	9.60	9.36	9.20	9.07	8.98	8.90
	.99	21.2	18.0	16.7	16.0	15.5	15.2	15.0	14.8	14.7
	.995	31.3	26.3	24.3	23.2	22.5	22.0	21.6	21.4	21.1
5	.50	0.528	0.799	0.907	0.965	1.00	1.02	1.04	1.05	1.06
	.90	4.06	3.78	3.62	3.52	3.45	3.40	3.37	3.34	3.32
	.95	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77
	.975	10.0	8.43	7.76	7.39	7.15	6.98	6.85	6.76	6.68
	.99	16.3	13.3	12.1	11.4	11.0	10.7	10.5	10.3	10.2
	.995	22.8	18.3	16.5	15.6	14.9	14.5	14.2	14.0	13.8
6	.50	0.515	0.780	0.886	0.942	0.977	1.00	1.02	1.03	1.04
	.90	3.78	3.46	3.29	3.18	3.11	3.05	3.01	2.98	2.96
	.95	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10
	.975	8.81	7.26	6.60	6.23	5.99	5.82	5.70	5.60	5.52
	.99	13.7	10.9	9.78	9.15	8.75	8.47	8.26	8.10	7.98
	.995	18.6	14.5	12.9	12.0	11.5	11.1	10.8	10.6	10.4
7	.50	0.506	0.767	0.871	0.926	0.960	0.983	1.00	1.01	1.02
	.90	3.59	3.26	3.07	2.96	2.88	2.83	2.78	2.75	2.72
	.95	5.59	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68
	.975	8.07	6.54	5.89	5.52	5.29	5.12	4.99	4.90	4.82
	.99	12.2	9.55	8.45	7.85	7.46	7.19	6.99	6.84	6.72
	.995	16.2	12.4	10.9	10.1	9.52	9.16	8.89	8.68	8.51

TABLE A.4 (continued) Percentiles of the F Distribution

Den. df	A	Numerator df								
		10	12	15	20	24	30	60	120	∞
1	.50	2.04	2.07	2.09	2.12	2.13	2.15	2.17	2.18	2.20
	.90	60.2	60.7	61.2	61.7	62.0	62.3	62.8	63.1	63.3
	.95	242	244	246	248	249	250	252	253	254
	.975	969	977	985	993	997	1,001	1,010	1,014	1,018
	.99	6,056	6,106	6,157	6,209	6,235	6,261	6,313	6,339	6,366
	.995	24,224	24,426	24,630	24,836	24,940	25,044	25,253	25,359	25,464
2	.50	1.34	1.36	1.38	1.39	1.40	1.41	1.43	1.43	1.44
	.90	9.39	9.41	9.42	9.44	9.45	9.46	9.47	9.48	9.49
	.95	19.4	19.4	19.4	19.4	19.5	19.5	19.5	19.5	19.5
	.975	39.4	39.4	39.4	39.4	39.5	39.5	39.5	39.5	39.5
	.99	99.4	99.4	99.4	99.4	99.5	99.5	99.5	99.5	99.5
	.995	199	199	199	199	199	199	199	199	200
3	.50	1.18	1.20	1.21	1.23	1.23	1.24	1.25	1.26	1.27
	.90	5.23	5.22	5.20	5.18	5.18	5.17	5.15	5.14	5.13
	.95	8.79	8.74	8.70	8.66	8.64	8.62	8.57	8.55	8.53
	.975	14.4	14.3	14.3	14.2	14.1	14.1	14.0	13.9	13.9
	.99	27.2	27.1	26.9	26.7	26.6	26.5	26.3	26.2	26.1
	.995	43.7	43.4	43.1	42.8	42.6	42.5	42.1	42.0	41.8
4	.50	1.11	1.13	1.14	1.15	1.16	1.16	1.18	1.18	1.19
	.90	3.92	3.90	3.87	3.84	3.83	3.82	3.79	3.78	3.76
	.95	5.96	5.91	5.86	5.80	5.77	5.75	5.69	5.66	5.63
	.975	8.84	8.75	8.66	8.56	8.51	8.46	8.36	8.31	8.26
	.99	14.5	14.4	14.2	14.0	13.9	13.8	13.7	13.6	13.5
	.995	21.0	20.7	20.4	20.2	20.0	19.9	19.6	19.5	19.3
5	.50	1.07	1.09	1.10	1.11	1.12	1.12	1.14	1.14	1.15
	.90	3.30	3.27	3.24	3.21	3.19	3.17	3.14	3.12	3.11
	.95	4.74	4.68	4.62	4.56	4.53	4.50	4.43	4.40	4.37
	.975	6.62	6.52	6.43	6.33	6.28	6.23	6.12	6.07	6.02
	.99	10.1	9.89	9.72	9.55	9.47	9.38	9.20	9.11	9.02
	.995	13.6	13.4	13.1	12.9	12.8	12.7	12.4	12.3	12.1
6	.50	1.05	1.06	1.07	1.08	1.09	1.10	1.11	1.12	1.12
	.90	2.94	2.90	2.87	2.84	2.82	2.80	2.76	2.74	2.72
	.95	4.06	4.00	3.94	3.87	3.84	3.81	3.74	3.70	3.67
	.975	5.46	5.37	5.27	5.17	5.12	5.07	4.96	4.90	4.85
	.99	7.87	7.72	7.56	7.40	7.31	7.23	7.06	6.97	6.88
	.995	10.2	10.0	9.81	9.59	9.47	9.36	9.12	9.00	8.88
7	.50	1.03	1.04	1.05	1.07	1.07	1.08	1.09	1.10	1.10
	.90	2.70	2.67	2.63	2.59	2.58	2.56	2.51	2.49	2.47
	.95	3.64	3.57	3.51	3.44	3.41	3.38	3.30	3.27	3.23
	.975	4.76	4.67	4.57	4.47	4.42	4.36	4.25	4.20	4.14
	.99	6.62	6.51	6.36	6.16	6.07	5.99	5.82	5.74	5.65
	.995	8.38	8.18	7.97	7.75	7.65	7.53	7.31	7.19	7.08

TABLE A.4 (continued) Percentiles of the F Distribution

Den. df	4	Numerator df									
		1	2	3	4	5	6	7	8	9	∞
8	.50	0.499	0.757	0.860	0.915	0.948	0.971	0.988	1.00	1.01	1.09
	.90	3.46	3.11	2.92	2.81	2.73	2.67	2.62	2.59	2.56	2.29
	.95	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39	2.93
	.975	7.57	6.06	5.42	5.05	4.82	4.65	4.53	4.43	4.36	3.67
	.99	11.3	8.65	7.59	7.01	6.63	6.37	6.18	6.03	5.91	4.86
9	.50	0.494	0.749	0.852	0.906	0.939	0.962	0.978	0.990	1.00	1.08
	.90	3.36	3.01	2.81	2.69	2.61	2.55	2.51	2.47	2.44	2.16
	.95	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18	2.71
	.975	7.21	5.71	5.08	4.72	4.48	4.32	4.20	4.10	4.03	3.33
	.99	10.6	8.02	6.99	6.42	6.06	5.80	5.61	5.47	5.35	4.31
10	.50	0.490	0.743	0.845	0.899	0.932	0.954	0.971	0.983	0.992	1.07
	.90	3.29	2.92	2.73	2.61	2.52	2.46	2.41	2.38	2.35	2.06
	.95	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02	2.54
	.975	6.94	5.46	4.83	4.47	4.24	4.07	3.95	3.85	3.78	3.08
	.99	10.0	7.36	6.35	5.99	5.64	5.39	5.20	5.06	4.94	3.91
12	.50	0.484	0.735	0.835	0.888	0.921	0.943	0.959	0.972	0.981	1.06
	.90	3.18	2.81	2.61	2.48	2.39	2.33	2.28	2.24	2.21	1.90
	.95	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80	2.30
	.975	6.55	5.10	4.47	4.12	3.89	3.73	3.61	3.51	3.44	2.72
	.99	9.33	6.93	5.95	5.41	5.06	4.82	4.64	4.50	4.39	3.36
15	.50	0.478	0.726	0.826	0.878	0.911	0.933	0.949	0.960	0.970	1.05
	.90	3.07	2.70	2.49	2.36	2.27	2.21	2.16	2.12	2.09	1.76
	.95	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59	2.07
	.975	6.20	4.77	4.15	3.80	3.58	3.41	3.29	3.20	3.12	2.42
	.99	8.68	6.36	5.42	4.89	4.56	4.32	4.14	4.00	3.89	2.87
20	.50	0.472	0.718	0.816	0.868	0.900	0.922	0.938	0.950	0.959	1.03
	.90	2.97	2.59	2.38	2.25	2.16	2.09	2.04	2.00	1.96	1.61
	.95	4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39	1.84
	.975	5.87	4.46	3.86	3.51	3.29	3.13	3.01	2.91	2.84	2.09
	.99	8.10	5.85	4.94	4.43	4.10	3.87	3.70	3.56	3.46	2.42
24	.50	0.469	0.714	0.812	0.863	0.895	0.917	0.932	0.944	0.953	1.03
	.90	2.93	2.54	2.33	2.19	2.10	2.04	1.98	1.94	1.91	1.53
	.95	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30	1.73
	.975	5.72	4.32	3.72	3.38	3.15	2.99	2.87	2.78	2.70	1.94
	.99	7.82	5.61	4.72	4.22	3.90	3.67	3.50	3.36	3.26	2.21

TABLE A.4 (continued) Percentiles of the F Distribution

Den. df	4	Numerator df									
		10	12	15	20	24	30	60	120	∞	∞
8	.50	1.02	1.03	1.04	1.05	1.06	1.07	1.08	1.08	1.08	1.09
	.90	2.54	2.50	2.46	2.42	2.40	2.38	2.34	2.32	2.32	2.29
	.95	3.35	3.28	3.22	3.15	3.12	3.08	3.01	2.97	2.97	2.93
	.975	4.30	4.20	4.10	4.00	3.95	3.89	3.78	3.73	3.73	3.67
	.99	5.81	5.67	5.52	5.36	5.28	5.20	5.03	4.95	4.95	4.86
9	.50	1.01	1.02	1.03	1.04	1.05	1.05	1.07	1.07	1.07	1.08
	.90	2.42	2.38	2.34	2.30	2.28	2.25	2.21	2.18	2.18	2.16
	.95	3.14	3.07	3.01	2.94	2.90	2.86	2.79	2.75	2.75	2.71
	.975	3.96	3.87	3.77	3.67	3.61	3.56	3.45	3.39	3.39	3.33
	.99	5.26	5.11	4.96	4.81	4.73	4.65	4.48	4.40	4.40	4.31
10	.50	1.00	1.01	1.02	1.03	1.04	1.05	1.06	1.06	1.06	1.07
	.90	2.32	2.28	2.24	2.20	2.18	2.16	2.11	2.08	2.08	2.06
	.95	2.98	2.91	2.84	2.77	2.74	2.70	2.62	2.58	2.58	2.54
	.975	3.72	3.62	3.52	3.42	3.37	3.31	3.20	3.14	3.14	3.08
	.99	4.85	4.71	4.56	4.41	4.33	4.25	4.08	4.00	4.00	3.91
12	.50	0.989	1.00	1.01	1.02	1.03	1.03	1.05	1.05	1.05	1.06
	.90	2.19	2.15	2.10	2.06	2.04	2.01	1.96	1.93	1.93	1.90
	.95	2.75	2.69	2.62	2.54	2.51	2.47	2.38	2.34	2.34	2.30
	.975	3.37	3.28	3.18	3.07	3.02	2.96	2.85	2.79	2.79	2.72
	.99	4.30	4.16	4.01	3.86	3.78	3.70	3.54	3.45	3.45	3.36
15	.50	0.977	0.989	1.00	1.01	1.02	1.02	1.03	1.03	1.03	1.05
	.90	2.06	2.02	1.97	1.92	1.90	1.87	1.82	1.79	1.79	1.76
	.95	2.54	2.48	2.40	2.33	2.29	2.25	2.16	2.11	2.11	2.07
	.975	3.06	2.96	2.86	2.76	2.70	2.64	2.52	2.46	2.46	2.40
	.99	3.80	3.67	3.52	3.37	3.29	3.21	3.05	2.96	2.96	2.87
20	.50	0.966	0.977	0.989	1.00	1.01	1.01	1.02	1.02	1.02	1.03
	.90	1.94	1.89	1.84	1.79	1.77	1.74	1.68	1.64	1.64	1.61
	.95	2.35	2.28	2.20	2.12	2.08	2.04	1.95	1.90	1.90	1.84
	.975	2.77	2.68	2.57	2.46	2.41	2.35	2.22	2.16	2.16	2.09
	.99	3.37	3.23	3.09	2.94	2.86	2.78	2.61	2.52	2.52	2.42
24	.50	0.961	0.972	0.983	0.994	1.00	1.01	1.02	1.02	1.02	1.03
	.90	1.88	1.83	1.78	1.73	1.70	1.67	1.61	1.57	1.57	1.53
	.95	2.25	2.18	2.11	2.03	1.98	1.94	1.84	1.79	1.79	1.73
	.975	2.64	2.54	2.44	2.33	2.27	2.21	2.08	2.01	2.01	1.94
	.99	3.17	3.03	2.89	2.74	2.66	2.58	2.40	2.31	2.31	2.21

TABLE A.4 (continued) Percentiles of the F Distribution

Den. df	A	Numerator df									
		1	2	3	4	5	6	7	8	9	
30	.50	0.466	0.709	0.807	0.858	0.890	0.912	0.927	0.939	0.948	
	.90	2.88	2.49	2.28	2.14	2.05	1.98	1.93	1.88	1.85	
	.95	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21	
	.975	5.37	4.18	3.59	3.25	3.03	2.87	2.75	2.65	2.57	
	.99	7.56	5.39	4.51	4.02	3.70	3.47	3.30	3.17	3.07	
	.995	9.18	6.35	5.24	4.62	4.23	3.95	3.74	3.58	3.45	
	.999	13.3	8.77	7.05	6.12	5.53	5.12	4.82	4.58	4.39	
60	.50	0.461	0.701	0.798	0.849	0.880	0.901	0.917	0.928	0.937	
	.90	2.79	2.39	2.18	2.04	1.95	1.87	1.82	1.77	1.74	
	.95	4.00	3.15	2.76	2.53	2.37	2.25	2.17	2.10	2.04	
	.975	5.29	3.93	3.34	3.01	2.79					

TABLE A.4 (concluded) Percentiles of the F Distribution

Den. dfr	Numerator dfr											
	10	12	15	20	24	30	60	120	∞			
30	0.955	0.966	0.978	0.989	0.994	1.00	1.01	1.02	1.02			
.50	1.82	1.77	1.72	1.67	1.64	1.61	1.54	1.50	1.46			
.95	2.16	2.09	2.01	1.93	1.89	1.84	1.74	1.68	1.62			
.975	2.51	2.41	2.31	2.20	2.14	2.07	1.94	1.87	1.79			
.99	2.98	2.84	2.70	2.55	2.47	2.39	2.21	2.11	2.01			
.995	3.34	3.18	3.01	2.82	2.73	2.63	2.42	2.30	2.18			
.999	4.24	4.00	3.75	3.49	3.36	3.22	2.92	2.76	2.59			
60	0.945	0.956	0.967	0.978	0.983	0.989	1.00	1.01	1.01			
.50	1.71	1.66	1.60	1.54	1.51	1.48	1.40	1.35	1.29			
.95	1.99	1.92	1.84	1.75	1.70	1.65	1.53	1.47	1.39			
.975	2.27	2.17	2.06	1.94	1.88	1.82	1.67	1.58	1.48			
.99	2.63	2.50	2.35	2.20	2.12	2.03	1.84	1.73	1.60			
.995	2.90											

Source: Reprinted from Table 5 of Pearson and Hartley, *Biometrika Tables for Statisticians*, Volume 2, 1972, published by the Cambridge University Press, on behalf of The Biometrika Society, by permission of the authors and publishers.