

DEPARTMENT OF STATISTICS AND DEMOGRAPHY

SUPPLEMENTARY EXAMINATION, 2009/10

COURSE TITLE: INFERENTIAL STATISTICS I

COURSE CODE: ST 232

TIME ALLOWED: TWO (2) HOURS

INSTRUCTION: ANSWER ANY THREE QUESTIONS
ALL QUESTIONS CARRY EQUAL MARKS (20 MARKS)

SPECIAL REQUIREMENTS: SCIENTIFIC CALCULATORS AND STATISTICAL TABLES

**DO NOT OPEN THIS PAGE UNTIL PERMISSION HAS BEEN GRANTED BY THE
INVIGILATOR**

Question 1

- (a) A certain machine makes electrical resistors having a mean of 40 ohms and a standard deviation of 2 ohms. Assuming that the resistance follows a normal distribution, what percentage of resistors will have a resistance exceeding 43 ohms? **(6 marks)**
- (b) Given a normal distribution with $\mu=40$ and $\sigma=6$, find the value of 'x' that has (i) 45% of the area to the left and (ii) 14% of the area to the right. **(6 marks)**
- (c) The probability that a patient recovers from a rare disease is 0.4. If 100 people are known to have contracted this disease, what is the probability that less than 30 survive? Use the Normal approximation to the Binomial to answer this question. **(8 marks)**

Question 2

The manufacturer of a gasoline additive claims that the use of such additive increases gasoline mileage. A random sample of six cars was selected and these cars were driven for one week without the gasoline additive and then one week with the gasoline additive. The following table gives the miles per gallon for these cars without and with gasoline additive.

Without	24.6	28.3	18.9	23.7	15.4	29.5
With	26.3	31.7	18.2	25.3	18.3	30.9

- (a) Using 2.5% significance level, can you conclude that the use of gasoline additive increases the gasoline mileage? **(12 marks)**
- (b) Construct a 99 percent confidence interval for μ_d , where $\mu_d = \mu_{\text{without}} - \mu_{\text{with additive}}$ **(8 marks)**

Question 3

(a) A manufacturer claims that the average tensile strength of thread A exceeds the average tensile strength of thread B by at least 12 kilograms. To test his claim, 50 pieces of each type of thread are tested under similar conditions. Type A thread had an average tensile strength of 86.7 kilograms with a standard deviation of 6.28 kilograms while type B thread had an average tensile strength of 77.8 kilograms with a standard deviation of 5.61 kilograms. Test the manufacturer's claim using a 0.05 level of significance.

(12 marks)

(b) In a study to test whether or not there is a difference between the average heights of adult females in two different countries, random samples of size $n_1=120$ and $n_2=150$ yielded $\bar{x}_1 = 62.7inches$ and $\bar{x}_2 = 61.8inches$. Exclusive studies of similar kind have shown that it is reasonable to let $\sigma_1 = 2.50inches$ and $\sigma_2 = 2.62inches$. Test at 0.05 level of significance whether the difference between these two sample means is significant.

(8 marks)**Question 4**

A survey was conducted among 400 "affluent" Americans with household incomes of \$750, 000 or more per year to examine the relationship between the use of a financial advisor and ownership of stocks. The results of the survey are reflected in the following table:

		Own Stocks	Do Not Own Stocks
Use financial advisor	Yes	165	135
	No	43	57

At 5% significance level, can you conclude that the use of a financial advisor is independent of stock ownership for all affluent Americans?

(20 marks)

Question 5

- (a) A large chain retailer purchases a certain kind of electronic device from a manufacturer. The manufacturer indicates that the defective rate of the device is 3%. The inspector of the retailer randomly picks 20 items from the shipment. What is the probability that there will be at least one defective item among these 20? **(5 marks)**
- (b) An electrical firm manufactures light bulbs that have a length of life that is approximately normally distributed, with mean equal to 800 hours and a standard deviation of 40 hours. Find the probability that a random sample of 16 bulbs will have an average life of less than 775 hours. **(5marks)**
- (c) The average zinc concentration recovered from a sample of zinc measurements in 36 different locations is found to be 2.6 grams per millilitre. Find the 95% confidence interval for the mean zinc concentration in the river. Assume that the population standard deviation is 0.3. **(5 marks)**
- (d) Among 5000 marriage license applications chosen at random in a given year, there were 48 in which the woman was at least one year older than the man and among 400 marriage license applications chosen at random six years later, there were 68 in which the woman was at least one year older than the man. Construct a 99% confidence interval for the difference between the corresponding true proportions of marriage license applications in which the woman was at least one year older than the man. **(5 marks)**

END OF EXAM!!

Table E The Standard Normal Distribution										
z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359
0.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0753
0.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141
0.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517
0.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879
0.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224
0.6	.2257	.2291	.2324	.2357	.2389	.2422	.2454	.2486	.2517	.2549
0.7	.2580	.2611	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
0.8	.2881	.2910	.2939	.2967	.2995	.3023	.3051	.3078	.3106	.3133
0.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4265	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4625	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4960	.4961	.4962	.4963	.4964
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981
2.9	.4981	.4982	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986
3.0	.4987	.4987	.4987	.4988	.4988	.4989	.4989	.4989	.4990	.4990

Note: Use 0.4999 for z values above 3.09.

Source: Frederick Mosteller and Robert E. K. Rourke, *Sturdy Statistics*, Table A-1 (Reading, Mass.: Addison-Wesley, 1973). Reprinted with permission of the copyright owners.





Table IX Chi-Square Distribution Table


The entries in this table give the critical values of χ^2 for the specified number of degrees of freedom and areas in the right tail.



df	Area in the Right Tail under the Chi-square Distribution Curve									
	.995	.990	.975	.950	.900	.100	.050	.025	.010	.005
1	0.000	0.000	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.815	9.348	11.345	12.838
4	0.207	0.297	0.484	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.610	9.236	11.070	12.833	15.086	16.750
6	0.676	0.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	0.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278
8	1.344	1.646	2.180	2.733	3.490	13.362	15.507	17.535	20.090	21.955
9	1.735	2.088	2.700	3.325	4.168	14.684	16.919	19.023	21.666	23.589
10	2.156	2.558	3.247	3.940	4.865	15.987	18.307	20.483	23.209	25.188
11	2.603	3.053	3.816	4.575	5.578	17.275	19.675	21.920	24.725	26.757
12	3.074	3.571	4.404	5.226	6.304	18.549	21.026	23.337	26.217	28.300
13	3.565	4.107	5.009	5.892	7.042	19.812	22.362	24.736	27.688	29.819
14	4.075	4.660	5.629	6.571	7.790	21.064	23.685	26.119	29.141	31.319
15	4.601	5.229	6.262	7.261	8.547	22.307	24.996	27.488	30.578	32.801
16	5.142	5.812	6.908	7.962	9.312	23.542	26.296	28.845	32.000	34.267
17	5.697	6.408	7.564	8.672	10.085	24.769	27.587	30.191	33.409	35.718
18	6.265	7.015	8.231	9.390	10.865	25.989	28.869	31.526	34.805	37.156
19	6.844	7.633	8.907	10.117	11.651	27.204	30.144	32.852	36.191	38.582
20	7.434	8.260	9.591	10.851	12.443	28.412	31.410	34.170	37.566	39.997
21	8.034	8.897	10.283	11.591	13.240	29.615	32.671	35.479	38.932	41.401
22	8.643	9.542	10.982	12.338	14.041	30.813	33.924	36.781	40.289	42.796
23	9.260	10.196	11.689	13.091	14.848	32.007	35.172	38.076	41.638	44.181
24	9.886	10.856	12.401	13.848	15.659	33.196	36.415	39.364	42.980	45.559
25	10.520	11.524	13.120	14.611	16.473	34.382	37.652	40.646	44.314	46.928
26	11.160	12.198	13.844	15.379	17.292	35.563	38.885	41.923	45.642	48.290
27	11.808	12.879	14.573	16.151	18.114	36.741	40.113	43.195	46.963	49.645
28	12.461	13.565	15.308	16.928	18.939	37.916	41.337	44.461	48.278	50.993
29	13.121	14.256	16.047	17.708	19.768	39.087	42.557	45.722	49.588	52.336
30	13.787	14.953	16.791	18.493	20.599	40.256	43.773	46.979	50.892	53.672
40	20.707	22.164	24.433	26.509	29.051	51.805	55.758	59.342	63.691	66.766
50	27.991	29.707	32.357	34.764	37.689	63.167	67.505	71.420	76.154	79.490
60	35.534	37.485	40.482	43.188	46.459	74.397	79.082	83.298	88.379	91.952
70	43.275	45.442	48.758	51.739	55.329	85.527	90.531	95.023	100.425	104.215
80	51.172	53.540	57.153	60.391	64.278	96.578	101.879	106.629	112.329	116.321
90	59.196	61.754	65.647	69.126	73.291	107.565	113.145	118.136	124.116	128.299
100	67.328	70.065	74.222	77.929	82.358	118.498	124.342	129.561	135.807	140.169

Table VIII The t Distribution Table[†]

The entries in the table give the critical values of t for the specified number of degrees of freedom and areas in the right tail.



df	Area in the Right Tail under the t Distribution Curve					
	.10	.05	.025	.01	.005	.001
1	3.078	6.314	12.706	31.821	63.657	318.309
2	1.886	2.920	4.303	6.965	9.925	22.327
3	1.638	2.353	3.182	4.541	5.841	10.215
4	1.533	2.132	2.776	3.747	4.604	7.173
5	1.476	2.015	2.571	3.365	4.032	5.893
6	1.440	1.943	2.447	3.143	3.707	5.208
7	1.415	1.895	2.365	2.998	3.499	4.785
8	1.397	1.860	2.306	2.896	3.355	4.501
9	1.383	1.833	2.262	2.821	3.250	4.297
10	1.372	1.812	2.228	2.764	3.169	4.144
11	1.363	1.796	2.201	2.718	3.106	4.025
12	1.356	1.782	2.179	2.681	3.055	3.930
13	1.350	1.771	2.160	2.650	3.012	3.852
14	1.345	1.761	2.145	2.624	2.977	3.787
15	1.341	1.753	2.131	2.602	2.947	3.733
16	1.337	1.746	2.120	2.583	2.921	3.686
17	1.333	1.740	2.110	2.567	2.898	3.646
18	1.330	1.734	2.101	2.552	2.878	3.610
19	1.328	1.729	2.093	2.539	2.861	3.579
20	1.325	1.725	2.086	2.528	2.845	3.552
21	1.323	1.721	2.080	2.518	2.831	3.527
22	1.321	1.717	2.074	2.508	2.819	3.505
23	1.319	1.714	2.069	2.500	2.807	3.485
24	1.318	1.711	2.064	2.492	2.797	3.467
25	1.316	1.708	2.060	2.485	2.787	3.450
26	1.315	1.706	2.056	2.479	2.779	3.435
27	1.314	1.703	2.052	2.473	2.771	3.421
28	1.313	1.701	2.048	2.467	2.763	3.408
29	1.311	1.699	2.045	2.462	2.756	3.396
30	1.310	1.697	2.042	2.457	2.750	3.385
31	1.309	1.696	2.040	2.453	2.744	3.375
32	1.309	1.694	2.037	2.449	2.738	3.365
33	1.308	1.692	2.035	2.445	2.733	3.356
34	1.307	1.691	2.032	2.441	2.728	3.348
35	1.306	1.690	2.030	2.438	2.724	3.340
36	1.306	1.688	2.028	2.434	2.719	3.333
37	1.305	1.687	2.026	2.431	2.715	3.326
38	1.304	1.686	2.024	2.429	2.712	3.319
39	1.304	1.685	2.023	2.426	2.708	3.313
40	1.303	1.684	2.021	2.423	2.704	3.307
∞	1.282	1.645	1.960	2.326	2.576	3.090

[†]This table is an abbreviated version of Table VIII that appears in Appendix C. This table goes up to 40 degrees of freedom. For degrees of freedom from 41 to 70, use Table VIII of Appendix C.